- 1987;110:1617-30.
- 198/;110:1617-30.
   Lewis RA, Sumner AJ. The electrodiagnostic distinctions between chronic familial and acquired demyelinative neuropathies. *Neurology* 1982;32:592-6.
   Pollard JD. Chronic inflammatory demyelinating
- 103 Pollard JD. Chronic inflammatory demyelinating polyradiculoneuropathy. In: McLeod JG, ed. Inflammatory Neuropathy. Baillière's clinical neurology 1:1. London: Baillière Tindall, 1994:107-27.

  104 Kyle 1978. Monoclonal gammopathy of undetermined significance—natural history of 241 cases. Am J Med 1978;64:814-26.
- 1976;04:614-20.
  105 Kahn SN, Riches PG, Kohn J. Paraproteinaemia in neurological disease: incidence, association, and classification of monoclonal immunoglobulins. J Clin Pathol 1980;33:617-21.
- 1980;33:617-21.
   106 Bleasel AF, Hawke SHB, Pollard JD, McLeod JG. IgG monoclonal paraproteinaemia and peripheral neuropathy. J Neurol Neurosurg Psychiatry 1993;56:52-7.
   107 Thomas PK, Willison HJ. Paraproteinaemic neuropathy.
- In: McLeod JG, ed. Inflammatory neuropathies.

  Baillière's clinical neurology 3:1. London: Baillière

  Tindall, 1994:129-47.
- Tindall, 1994:129-47.

  108 Pestronk A, Cornblath DR, Ilyas AA, Baba H, Quarles RH, Griffin JW, et al. A treatable multifocal motor neuropathy with antibodies to GM1 ganglioside. Ann Neurol 1988;24:73-8.

  109 Pestronk A, Chaudhry V, Feldman EL, Griffin JW, Cornblath DR, Denys EH, et al. Lower motor neuron syndromes defined by patterns of weakness, nerve conduction abnormalities and high titres of antiglycolipid antibodies. Ann Neurol 1990;27:315-26. antibodies. Ann Neurol 1990;27:315-26.
- 110 Krarup C, Stewart MB, Sumner AJ, Pestronk A, Lipton SA. A syndrome of asymmetric limb weakness with motor conduction block. Neurology 1990;40:118-27.

- 111 Lange DJ, Trojaborg W, Latov N, Hays AP, Younger DS, Uncini A, et al. Multifocal motor neuropathy with conduction block: Is it a distinct clinical entity? Neurology 1992;42:497-505.
  112 Prineas JW. Pathology of inflammatory demyelinating neuropathies. In: McLeod JG, ed. Inflammatory neuropathies. Baillière's clinical neurology 3:1. London: Baillière Tindall, 1994:1-24.
- 113 Kaji R, Oka N, Tsuji T, Mezaki T, Nishio T, Akiguchi I, Kimura J. Pathological findings at the site of conduction block in multifocal motor neuropathy. Ann Neurol 1993;33:152-8.
- 114 Mellgren S, Conn DL, Stevens JC, Dyck PJ. Peripheral neuropathy in primary Sjögren's syndrome. Neurology 1989;39:390-4
- 1989;39:390-4.
  115 Gemignani F, Marbini A, Pavesi G, Di Vittorio S, Managelli P, Cenacchi G, Mancia D. Peripheral neuropathy associated with primary Sjögren's syndrome. *J Neurol Neurosurg Psychiatry* 1994;57:983-6.
  116 Griffin JW, Comblath DR, Alexander E. Ataxic sensory
- 116 Griffin JW, Cornblath DR, Alexander E. Ataxic sensory neuropathy and dorsal root ganglionitis associated with Sjögren's syndrome. Ann Neurol 1990;27:304-15.
  117 Said G. Inflammatory neuropathies associated with known infections (HIV, leprosy, Chagas' disease, Lyme disease). In: McLeod JG, ed. Inflammatory neuropathies. Baillière's clinical neurology 3:1. London: Baillière Tindall, 1994:149-72.
  118 So YT, Olney RK. Acute lumbosacral polyradiculoneuropathy in acquired immunodeficiency syndrome.
- ropathy in acquired immunodeficiency syndrome: Experience in 23 patients. Ann Neurol 1994;35:53–8. iabin TD, Swift TR, Jacobson RR. Leprosy. In: Dyck PJ, Thomas PK, Griffin JW, Low PA, Poduslo AN, eds. Peripheral neuropathy. 3rd ed. Philadelphia: WB Saunders, 1993:1354–79. 119 Sabin

## NEUROLOGICAL STAMP

## Atropa belladonna (deadly nightshade)

Atropa belladonna is a member of the nightshade family solanaceae. Its botanical name is derived from Atropos who, in Greek mythology, cut the thread of life. The red sap enlarges the pupils of the eye. The plant was named at the time of the Italian Renaissance in the days of the infamous Borgia family, when the ready availability of poison was popular. The ladies of the court used one of these poisons, belladonna, not to kill their rivals but to beautify and enlarge their pupils.

The solanaceous alkaloids are among the earliest effective treatments for parkinsonism. Belladonna contains atropine, scopolamine, and hyoscyamine. The ripe berries are sweet tasting and poisonous and attractive to children.

A medicinal plant series published and produced by Yugoslavia in 1965 shows the belladonna flower (Stanley Gibbons 1161, Scott 775).

L F HAAS

