

Variability of topsoil properties at the southern coast of Finland and the number of soil samples needed for the estimation of soil properties

RAILI JOKINEN

University of Helsinki, Department of Agricultural Chemistry, Sf-00710 Helsinki 71, Finland

Abstract: A total of 430 topsoil samples were collected from ten fields of the Viikki Experimental Farm, University of Helsinki. Particle size distribution, organic carbon content, pH(CaCl₂), exchangeable Ca, Mg, K contents, plant available P (Bray 1), 1 M KCl extractable (Al+H) content and effective cation exchange capacity of the soils were determined.

The coefficient of variation was used as indicator of the variability of soil properties within each field. The lowest coefficients of variation were observed for pH(CaCl₂) and the highest for exchangeable Mg 1 M KCl extractable (Al+H) and effective cation exchange capacity.

The results indicate that from 1 (pH(CaCl₂)) to 33 (exchangeable Mg) samples per hectare are needed from individual fields for strict level of accuracy in estimation of the soil properties. For determination of soil type (according to clay content) and organic carbon content on average 8 samples, and for the plant available P (Bray 1) and exchangeable Mg and K contents 10 to 16 samples per hectare appear sufficient. Four samples suffice for a less stringent, lax accurate determination of all properties.

The variability of soil properties is discussed from the viewpoint of agricultural advisory work and field experiments for agricultural research.

Introduction

As early as 1935 KIVINEN drew attention to the great variability in the chemical properties of soil even within the space of 30 or 40 metres. This he found to cause wide variations in the yields of the reference variety in a field experiment. KAILA and RYTI (1951) studied soil samples taken at distances of 2 metres and distances of 25 centimetres, concluding that it is difficult to obtain really representative samples for the estimation of soil properties. In a review article BECKETT and WEBSTER (1971) noted about 80 studies dealing with the variability of the properties of agricultural and forest soils. The subject has since been re-investigated in connection, for example, with regulation of nitrogen fertilization of farmland (LINDÉN 1979), forest management (QUESNEL and LAVKULICH 1980) and prediction of timber yields (BLYTH and MACLEOD 1978).

The aim of this study was to clarify the variability in soil properties as these may affect the interpretation of results obtained in field experiments and the recommendation made in the course of agricultural advisory work. The number of soil samples needed for accurate estimation of the properties is calculated.

Materials and methods

The material of the study consists of topsoil samples collected from the agricultural area of the Viikki Experimental Farm (University of Helsinki). This farm is situated near the Gulf of Finland. The geography of the fields is rather flat. In the soil profiles there appears strata with considerable differences in the particle size distribution originating in the time of deposition.

After harvestings of the crops, in September 1979 and 1981, the fields were marked with lines 40 metres apart, along which soil samples (volume 2 litres) representing the plough layer were taken at 40-metre intervals. Ten fields were sampled as follows:

Field number	Area ha	Number of samples	Samples/ha
49	7.27	45	6.2
54	17.18	99	5.8
84	8.21	42	5.1
86	7.37	46	6.2
88	5.22	34	6.5
89	5.28	31	5.9
94	1.08	6	5.6
96	10.29	66	6.4
97	3.60	23	6.4
98	5.82	38	6.5
Total	71.32	430	

About one third of the Viikki area of 232 ha was sampled (Fig. 1). For the minor part the fields number 54, 84 and 86 were at the time under fields experiments, no fertilization experiments.

The samples were kept on laboratory tables until they reached air-dry state, after which they were crushed to pass through a 2 mm-sieve. The properties of the soils were determined by the following methods:

- pH in 0.01 M CaCl₂ suspension, soil to solution ratio 1:2.5 (v/v), equilibration time 4 hours. The material was classified into four groups, where the pH(CaCl₂) value was 3.5-4.4, 4.5-5.4, 5.5-6.4 and >6.5.
- organic carbon, by wet combustion with K₂Cr₂O₇ and H₂SO₄ (conc.) and thereafter colorimetric determination. On the basis of the organic carbon content the mineral soils were classified into three groups of 1.7-3.4 %, 3.5-6.9 %, 7.0-11.5 %. The soils with organic carbon content 11.6-23.2 % are mull soils.
- particle sizes were determined by pipette method (ELONEN 1971). Soils where the clay (< 2 μm) content is less than 30 % are non-clay soils. In the total material there were 179 samples of fine sand soils, 21 samples of finer fine sand, 226 samples of sandy clay soils. The number of soil samples in the different groups defined by clay content, pH(CaCl₂) and organic carbon content are presented in Table 1.




Fig. 1. Field layout of the Viikki Experimental Farm, University of Helsinki.

- the exchangeable cations were extracted with 1 M neutral ammonium acetate. Calcium and magnesium were determined by atomic absorption spectrophotometry (Varian Techtron 1 00), with interference Sr and exchangeable K by flame photometry (Lange).
- the exchange acidity (Al+H) was displaced with 1 M KCl and titrated with 0.01 M NaOH.
- the effective cation exchange capacity (ECEC) was determined as the sum of (Ca+Mg) and (Al+H) extractable in 1 M KCl (KAILA 1971).
- the plant available form of soil phosphorus was extracted with 0.03 M NaF+0.025 M HCl by the Bray 1 test (BRAY and KURTZ 1945) modified by KAILA (1965) and determined by molybdenum blue method.

The mean (\bar{x}), standard deviation (s) and coefficient of variation in per cent (v) of the soil properties were calculated for each of the ten fields. The number of samples needed for accurate determination of the soil properties was calculated according to SNEDECOR (1948) by the equation $n = \frac{t^2 v^2}{p^2}$, where n = number of samples needed, t^2 = square of Students t , v^2 = square of coefficient of variation, and p = allowable error in per cent. The number of samples needed was calculated both with $t=5\%$, $p=10\%$ (strict level of accuracy = n_1) and with $t=10\%$, $p=25\%$ (lax level of accuracy = n_2).

Results

Except in field 54 only one crop was grown in each field, so that fertilization was the same over the whole field. On field 54 two different crops were grown, in both sampling years. Within a field main reason for the variation should then be the inherent heterogeneity of soil properties and the cultivation history.

Table 1 The number of soil samples in different classes of clay content, pH(CaCl₂) and organic carbon content in ten fields.

	Number of soil samples									
	Field number									
	49	54	84	86	88	89	94	96	97	98
Clay, %										
< 30	38	83	19	17	17	4		2	7	13
30-60	7	15	23	26	17	27	6	64	16	25
> 60		1		3						
Org. C, %										
1.7- 3.4	18	39	3	21	11			1	3	15
3.5- 6.9	22	47	39	25	20	7	2	28	9	21
7.0-11.5	2	13			3	23	3	37	11	2
11.6-23.2	3					1	1			
pH(CaCl ₂)										
3.5- 4.4					3		1	1	7	3
4.5- 5.4	7	41	29	14	30	27	5	52	16	22
5.5- 6.4	29	54	12	32	1	4		12		12
6.5-	9	4	1					1		1

The coefficient of variation of pH(CaCl₂) was between 5.4 and 10 % in the individual fields (Table 2). The difference between maximum and minimum pH(CaCl₂) values in a single field was at the lowest 1.0 unit (field 97) and at the highest 2.3 units (field 98). The distance between the extreme values in field 98 was about 170 metres. In the field 49 the pH(CaCl₂) values for the two adjacent points (distance apart 40 metres) with greatest difference in value were 6.1 and 4.9. Were the pH(CaCl₂) of the soil to be adopted as the indicator of the liming requirement, the minimum, maximum and mean values would indicate the addition of three different amounts of liming agents.

Because the pH scale is logarithmic the coefficient of variation of pH(CaCl₂) is not comparable to the coefficients for other soil properties.

The range in the organic carbon content was widest in field 49 from 1.8 to 14.6 % (Table 3) and the coefficient of variation was highest there too (Table 2). As high carbon contents were observed in field 94 as in field 49, but the material was concentrated near the mean and the coefficient of variation remained low. The change in carbon content in field 49 occurred gradually, unlike pH(CaCl₂). The great differences in organic carbon content cause differences in the water and nutrient retention capacity of the soil and can lead to uneven maturity of the crops.

The range of the clay content was especially wide in fields 54 and 86 (Table 3). In field 54 the clay content changed within a distance of 100 metres from 11 to 65 % and the fine sand content from 76 to 17 %. In field 86 the distance between minimum and maximum clay contents (7 and 62 %) was

Table 2. The coefficient of variation (v %), the number of soil samples needed for the strict (n₁) and lax (n₂) level of accuracy in determination of soil properties in the individual fields, and the average values (field 94 omitted).

	v %	n ₁	n ₂	v %	n ₁	n ₂	v %	n ₁	n ₂	v %	n ₁	n ₂
	<u>< 2 μm, %</u>			<u>2-20 μm, %</u>			<u>20-200 μm, %</u>			<u>Org. C, %</u>		
49	49.5	14	1.5	43.8	11	1.2	16.9	2	0.2	62.2	22	2.4
54	52.4	6	0.7	32.2	2	0.3	21.9	1	0.1	40.0	4	0.4
84	21.2	2	0.2	21.9	2	0.3	19.8	2	0.2	17.4	2	0.2
86	42.7	10	1.1	29.5	5	0.5	33.1	6	0.7	16.7	2	0.2
88	40.5	13	1.4	30.4	7	0.8	25.8	5	0.6	37.8	11	1.3
89	20.1	3	0.4	16.5	2	0.2	21.4	4	0.4	22.2	4	0.4
94	16.9	16	1.6	26.9	40	0.4	12.5	9	0.9	38.8	99	8.4
96	11.2	1	0.1	11.9	1	0.1	16.3	1	0.1	18.6	1	0.1
97	30.6	11	1.2	34.5	14	1.6	44.8	24	2.6	32.8	13	1.4
98	34.0	8	0.9	22.0	3	0.4	32.3	7	0.8	39.0	11	1.2
Average	31.9	8.4	0.9	27.0	5.2	0.6	24.5	5.8	0.6	32.6	7.8	0.8
	<u>pH(CaCl₂)</u>			<u>Ca mg/kg</u>			<u>Mg mg/kg</u>			<u>K mg/kg</u>		
49	10.0	0.5	0.1	31.3	5	0.6	56.9	18	2.0	64.0	23	2.5
54	8.9	0.2	0.0	25.0	1	0.2	79.7	15	1.6	53.9	7	0.7
84	7.6	0.2	0.0	20.0	2	0.2	38.8	7	0.8	36.0	6	0.7
86	5.4	0.2	0.0	28.5	4	0.5	58.0	19	2.1	29.8	5	0.5
88	8.2	0.5	0.1	20.9	3	0.4	41.9	14	1.5	30.3	7	0.8
89	6.0	0.2	0.0	15.9	2	0.2	24.8	5	0.5	64.2	32	3.6
94	8.3	5.0	0.5	42.4	100	10.0	29.4	48	4.8	48.3	130	13.0
96	9.8	0.3	0.0	24.4	2	0.3	41.6	7	0.7	26.4	3	0.3
97	7.1	0.6	0.1	17.9	4	0.4	52.3	33	3.6	38.9	18	2.0
98	9.4	0.6	0.1	22.5	4	0.4	59.9	25	2.8	39.2	11	1.2
Average	8.1	0.4	0.1	24.9	3.0	0.4	48.3	15.9	1.7	43.1	12.4	1.4
	<u>P(Bray 1)mg/kg</u>			<u>(Al+H) me/kg</u>			<u>ECEC mg/kg</u>					
49	59.8	20	2.2	141.4	111	12.4	30.4	5	0.6			
54	38.6	3	0.4	68.3	11	1.2	25.7	2	0.2			
84	36.2	7	0.7	66.7	22	2.4	14.9	1	0.1			
86	41.2	9	1.0	33.3	6	0.7	30.7	5	0.6			
88	33.0	9	1.0	78.6	35	3.9	226.5	406	32.1			
89	28.9	7	0.7	60.0	28	3.1	13.8	1	0.2			
94	58.9	193	19.3	61.1	207	20.8	133.3	988	99.1			
96	50.9	10	1.1	50.9	10	1.1	72.7	21	2.3			
97	35.1	15	1.6	35.2	15	1.6	64.0	49	5.3			
98	42.8	13	1.4	43.0	13	1.4	122.2	105	11.6			
Average	42.5	10.3	1.1	63.9	27.9	3.1	73.4	158	15.2			

about 90 metres. The change in the soil type from fine sand to heavy clay cannot be without effect on the cation content of the soil and on the fertilizer requirement.

The coefficient of variation in particle size distribution, in the amounts of clay, silt or fine sand fractions, were below 50 % except for field 54 (Table 2). In field 96 the coefficient of variation in clay content was low (11%), since there sandy clay samples accounted for 64 out of the total 66 samples.

Table 3. The mean value, standard deviation ($\bar{x}\pm s$) and range for soil properties of individual fields.

	Field number										
	49	54	84	86	88	89	94	96	97	98	
Area, hectares	7.27	17.18	8.21	7.37	5.22	5.28	1.08	10.29	3.60	5.82	
Number of samples	45	99	42	46	34	31	6	66	23	38	
Samples, per hectare	6.2	5.8	5.1	6.2	6.5	5.9	5.6	6.4	6.4	6.5	
Particle-size analysis											
<2 μm , %	$\bar{x}\pm s$	20 \pm 10	21 \pm 11	30 \pm 6	36 \pm 15	28 \pm 11	38 \pm 8	42 \pm 7	42 \pm 5	33 \pm 10	33 \pm 11
	range	9-46	6-65	19-47	7-62	9-49	14-50	30-50	27-50	14-50	9-51
2-20 μm , %	$\bar{x}\pm s$	11 \pm 5	12 \pm 4	18 \pm 4	13 \pm 4	13 \pm 4	16 \pm 3	28 \pm 8	27 \pm 3	26 \pm 9	28 \pm 6
	range	6-35	3-24	4-27	6-22	6-20	7-21	23-43	16-34	8-37	15-40
20-200 μm , %	$\bar{x}\pm s$	63 \pm 11	63 \pm 14	48 \pm 10	47 \pm 16	56 \pm 15	41 \pm 9	27 \pm 3	28 \pm 5	36 \pm 16	36 \pm 12
	range	34-80	17-90	35-78	23-79	31-82	28-70	24-33	22-45	19-69	21-64
Organic C, %	$\bar{x}\pm s$	4.5 \pm 2.8	4.5 \pm 1.8	4.6 \pm 0.8	3.6 \pm 0.6	4.5 \pm 1.7	8.1 \pm 1.8	8.5 \pm 3.3	7.0 \pm 1.3	6.1 \pm 2.0	4.1 \pm 1.6
	range	1.8-14.6	2.2-11.2	2.2-5.9	2.1-4.8	2.5-9.4	3.5-11.7	5.2-14.4	2.6-9.5	2.5-9.2	2.3-8.5
pH(CaCl ₂)	$\bar{x}\pm s$	6.0 \pm 0.6	5.6 \pm 0.5	5.3 \pm 0.4	5.6 \pm 0.3	4.9 \pm 0.4	5.0 \pm 0.3	4.8 \pm 0.4	5.1 \pm 0.5	4.6 \pm 0.3	5.3 \pm 0.5
	range	4.6-6.8	4.5-6.6	4.8-6.6	4.9-6.1	4.2-5.6	4.5-5.6	4.0-5.1	4.4-6.5	4.2-5.2	4.3-6.6
P (Bray 1) mg/kg	$\bar{x}\pm s$	82 \pm 49	171 \pm 66	94 \pm 34	131 \pm 54	109 \pm 36	114 \pm 33	18 \pm 11	55 \pm 28	128 \pm 45	100 \pm 43
	range	5-206	53-355	44-196	64-292	50-176	53-194	2-30	14-151	63-228	32-158
Exchangeable (pH 7)											
Ca mg/kg	$\bar{x}\pm s$	2757 \pm 862	2228 \pm 557	2195 \pm 439	2312 \pm 658	1610 \pm 337	2760 \pm 439	2688 \pm 1139	3198 \pm 779	1593 \pm 285	1941 \pm 436
	range	1275-5845	1035-3975	1495-3930	750-3550	1000-2700	1775-3531	502-3537	1136-4780	1215-2269	950-3190
Mg mg/kg	$\bar{x}\pm s$	204 \pm 119	200 \pm 159	160 \pm 62	364 \pm 211	124 \pm 52	145 \pm 36	205 \pm 61	144 \pm 60	94 \pm 50	172 \pm 103
	range	44-585	69-708	84-373	55-810	48-261	73-224	137-298	70-357	45-277	36-598
K mg/kg	$\bar{x}\pm s$	198 \pm 127	306 \pm 164	436 \pm 157	356 \pm 109	234 \pm 71	246 \pm 158	296 \pm 143	276 \pm 73	357 \pm 139	355 \pm 139
	range	53-510	130-1440	168-838	210-700	129-374	114-970	174-350	155-480	125-765	65-617
Effective CEC											
me/kg	$\bar{x}\pm s$	134 \pm 41	117 \pm 30	117 \pm 18	140 \pm 43	98 \pm 222	152 \pm 21	162 \pm 34	173 \pm 31	105 \pm 19	114 \pm 23
	range	62-288	51-194	93-168	43-212	61-143	86-190	105-198	88-250	70-136	52-157
1 M KCl extract.											
(Al+H) me/kg	$\bar{x}\pm s$	2.9 \pm 4.1	4.1 \pm 2.8	5.4 \pm 3.6	3.0 \pm 0.8	14.4 \pm 10.9	9.9 \pm 6.0	17.9 \pm 24.0	10.8 \pm 7.6	25.1 \pm 15.6	9.3 \pm 10.7
	range	1.0-25.2	1.0-15.4	1.2-16.8	2.0-6.2	3.0-39.7	3.0-23.3	4.8-66.2	2.3-33.5	3.9-47.6	1.1-39.2

In the P (Bray 1) content, exchangeable Mg and K contents, (Al+H) content and ECEC the coefficients of variation were over 30 % for almost all fields. Exceptionally high coefficients of variation were recorded for ECEC in three fields. The range of these properties was not widest in the same field for which the greatest coefficients of variation were observed. In the fields where the coefficient of variation in the exchangeable Mg was high there was also high in clay content. The coefficient of variation in the exchangeable Ca remained below 30 % except in fields 49 (31 %) and 94 (42 %).

The number of soil samples per hectare needed to satisfy the strict (n_1) and lax (n_2) accuracy classes as defined above was calculated separately for each field. (Field 94 was omitted because of its small area). In general, n_1 for pH(CaCl₂) was less than one sample per hectare. For the determination of the soil type according to the clay content n_1 was between 2 and 16 samples and for n_2 about one sample.

For strict level of accuracy in determination of the organic carbon content the number of soil samples needed was between 2 and 22, and for lax level of accuracy about one sample per hectare.

Determinations of the exchangeable K and Mg contents with the strict criterion of accuracy demanded from 3 to 32 and from 5 to 33 samples per hectare, respectively. For the determination of the plant-available P content the number of soil samples demanded varied from 3 to 20 and for the

exchangeable Ca content from one to five with the strict criterion of accuracy. Lax accurate determination of the nutrient contents could be satisfied by a collection of one (Ca) to four (K and Mg) samples. Because both low and high coefficients of variations were found for 1 M KCl extractable (Al+H) and ECEC, the number of soil samples needed for lax accurate determination varied from 0.1 to 32 per hectare.

For the determination of soil type and organic carbon content with the strict accuracy criterion on average 8 soil samples per hectare was found necessary. Correspondingly the determinations of plant available P and exchangeable Mg and K demanded 10 and 16 samples, respectively.

Diskussion

The 430 topsoil samples collected from the Viikki Experimental Farm represented mainly mineral soils. The mean $\text{pH}(\text{CaCl}_2)$ was near the average value of Finnish mineral soils (SIPPOLA and TARES 1978), considering that $\text{pH}(\text{H}_2\text{O}) = 0.5 + \text{pH}(\text{CaCl}_2)$ (RYTI 1965). The exchangeable Ca and K contents were higher than the average values reported for Finnish mineral soils by KAILA (1973) and SIPPOLA and TARES (1978), while the exchangeable Mg content does not deviate from the values of the same authors. The ECEC's are likewise in good accordance with an earlier study (KAILA 1971). The plant-available P contents are higher than the contents reported by KAILA (1965) using the same method probably because of the heavy phosphorus fertilization in the sixties and seventies. The drilling of fertilizers may also cause some differences in the nutrient contents of soil samples taken between rows and along the rows. This effect was probably slight in the present study, however, since samples were taken in the autumn after the harvesting of the crops (URVAS and JUSSILA 1979).

The variability in soil properties was studied in ten fields, revealing a wide range in all the soil properties in individual fields. The ranges obtained by KAILA and RYTI (1951) within 100 square metres and within 1 square metre were slightly narrower than in this material within 1 to 17 hectares. The size of the sampling area apparently has little effect of the ranges found in the soil properties (HEMINGWAY 1955).

The coefficients of variation observed by BALL and WILLIAMS (1968) for uncultivated and unfertilized soils in North Wales were almost the same as in this study for cultivated and fertilized soils. Likewise they reported the highest coefficient of variation for the exchangeable cation contents.

For the present agricultural advisory work in Finland about 1.5 soil samples are collected per hectare (KURKI 1982). In this study one to two samples were found adequate for strict accurate determination only of $\text{pH}(\text{CaCl}_2)$. The number of soil samples should be decided according to the most variable property, which in this material was the exchangeable Mg content. Even for the lax accurate determination 4 soil samples per hectare

were needed. LINDÉN (1979) suggested collecting about 10 soil samples and for field experiments (LINDÉN 1981) 14 cores per plot (108 square metres).

When field experiments are being laid out the determination of soil properties in advance is important, and the number of soil samples should rather be too high than too low. Several soil samples, each collected from a different sampling point, are more informative than a single sample made up of subsamples from different sampling points. Too little attention has thus far been paid to the density and mode of sampling.

References

- BALL, D. F. & WILLIAMS, W. M. 1968. Variability of soil chemical properties in two uncultivated brown earths. *J. Soil Sci.* 19: 379–391.
- BECKETT, P. H. T. & WEBSTER, R. 1971. Soil variability: A review. *Soils Fertil.* 34: 1–15.
- BLYTH, J. F. & MACLEOD, D. A. 1978. The significance of soil variability for forest soil studies in north-east Scotland. *J. Soil Sci.* 29: 419–430.
- BRAY, R. H. & KURTZ, L. T. 1945. Determination of total, organic and available forms of phosphorus in soils. *Soil Sci.* 59: 39–45.
- ELONEN, P. 1971. Particle-size analysis. *Acta Agric. Fenn.* 122: 1–122.
- HEMINGWAY, R. G. 1955. Soil sampling errors and advisory analysis. *J. Agric. Sci.* 46: 1–8.
- KAILA, A. 1965. Some phosphorus test values and fractions of inorganic phosphorus in soils. *J. Scient. Agric. Soc. Finl.* 37: 175–185.
- 1971. Effective cation-exchange capacity in Finnish mineral soils. *J. Scient. Agric. Soc. Finl.* 43: 178–186.
- 1973. Calcium, magnesium and potassium in mineral soils from southern half of Finland. *J. Scient. Agric. Soc. Finl.* 45: 254–261.
- & RYTI, R. 1951. Observations on factors influencing the results of chemical soil tests. *Acta Agric. Scand.* 1: 271–281.
- KIVINEN, E. 1935. Über die Ungleichmässigkeit des Ackerbodens. *J. Scient. Agric. Soc. Finl.* 7: 1–17.
- KURKI, M. 1982. Suomen peltojen viljavuudesta III. Summary: On the fertility of Finnish tilled fields in the light of investigations of soil fertility carried out in the years 1955–1980. 181 p. Helsinki.
- LINDÉN, B. 1979. Kvävegödsling baserad på bestämning av mineralkväveförrådet i marken. Längesrapport om N-prognosverksamhet i några europeiska länder och i Nordamerika. Summary: Nitrogen fertilization recommendations based on determination of mineral nitrogen in soils. Research and extension facilities for N-prognosis in some European countries and in North America. Swedish Univ. of Agric. Sci., Dept. Soil Sci. Div. Soil Fert. Rep. 122: 1–80, 8 Tabl., 22 fig., 3 enclosures.
- 1981. Ammonium- och nitratkvävet rörelser och fördelning i marken II. Metoder för mineralkväveprovtagning och -analyser. Summary: Movement and distribution of ammonium- and nitrate-N in the soil II. Methods of sampling and analysing mineral nitrogen. Swedish Univ. of Agric. Sci., Dept. Soil Sci., Div. Soil Fert. Rep. 137: 1–79, 8 Tabl., 32 Fig.
- QUESNEL, H. J. & LAVKULICH, L. M. 1980. Nutrient variability of forest floor near Port Hardy, British Columbia, Canada. *Can. J. Soil Sci.* 60: 565–573.
- RYTI, R. 1965. On the determination of soil pH. *J. Scient. Agric. Soc. Finl.* 37: 51–60.
- SIPPOLA, J. & TARES, T. 1978. The soluble content of mineral elements in cultivated Finnish soils. *Acta Agric. Scand. Suppl.* 20: 11–25.
- SNEDECOR, G. W. 1948. *Statistical methods*. 485 p. 4th Ed., 3rd Print. Ames Iowa.
- URVAS, L. & JUSSILA, M. 1979. Rivilannoituksen vaikutus viljavuustutkimuksen tulokseen. Maatalouden tutkimuskeskus, Maantutkimuslaitoksen Tiedote 7: 1–10.

Muokkauskerroksessa ominaisuuksien vaihtelevuus Suomenlahden rannikon pelloilla ja analyysejä varten tarvittavien näytteiden lukumäärä

Raili Jokinen

Helsingin yliopisto, Maanviljelyskemian laitos, 00710 Helsinki 71

Tutkimusta varten kerätiin syksyllä 1979 ja 1981 Helsingin yliopiston Viikin koetilan pelloilta kymmeltä lohkolta yhteensä 430 muokkauskerroksen näytettä. Lohkoille merkittiin näytteiden ottolinjat 40 m välein ja kullekin linjalle näytteiden ottoapaikat 40 m välein. Hehtaaria kohti otettujen näytteiden määrä vaihteli 5.1–6.5.

Näytteistä analysoitiin raekoostumus, orgaanisen hiilen pitoisuus, $\text{pH}(\text{CaCl}_2)$, kasveille käyttökelpoinen fosfori (Bray 1 testillä), neutraalin ammoniumasetaatin vaihtamat kalsium, magnesium ja kalium, efektiivinen kationinvaihtokapasiteetti sekä 1 M kaliumkloridiin uuttuvan aluminiumin ja vedyn summa. Ominaisuuksien vaihtelevuutta eri lohkojen välillä ja lohkojen sisällä tutkittiin variaatiokertoimen avulla. Analyysejä varten tarvittavien näytteiden lukumäärä laskettiin kahdella tulosten tarkkuuden tasolla: n_1 = ankara tarkkuus (sallittu poikkeama 10 %) ja n_2 = kohtalainen tarkkuus (sallittu poikkeama 25 %).

Kullakin lohkolta $\text{pH}(\text{CaCl}_2)$ -luvun variaatiokertoimet olivat alhaiset (alle 10 %), vaikka $\text{pH}(\text{CaCl}_2)$ saattoi lohkon sisällä vaihdella 4.6–6.8 (taulukko 2 ja 3). Suurimmat kertoimen arvot yksittäisillä lohkoilla todettiin vaihtuvan magnesiumin pitoisuuden, efektiivisen kationinvaihtokapasiteetin ja 1 M kaliumkloridiin uuttuvan aluminiumin ja vedyn summan analyysituloksissa. Eri ominaisuuksien vaihteluväli lohkojen sisällä oli laaja (taulukko 3).

Otettaessa maanäytteitä kenttäkokeita tai neuvontaa varten näytteiden lukumäärä tulisi ratkaista eniten vaihtelevan kulloinkin kyseeseen tulevan maan ominaisuuden perusteella. Näyttää siltä, että keskimäärin kahdeksan näytettä hehtaarilta tarvittaisiin maalajin (savenpitoisuus) ja orgaanisen hiilen pitoisuuden määrittämiseen, sekä 10–16 näytettä kasveille käyttökelpoisen fosforin tai vaihtuvien kationien pitoisuuden määrittämiseen ankarat vaatimukset täyttävällä tarkkuudella. Kohtalaisen tarkkuuden täyttävät analyysitulokset saataneen noin neljästä näytteestä hehtaarilta.

Tässä tutkimuksessa saadut näytteiden lukumäärää koskevat tulokset soveltunevat paremmin neuvonnan kuin tutkimuksen tarkoituksiin. Kenttäkokeista maanäytteitä tulisi ottaa hehtaaria kohti huomattavasti enemmän kuin edellä esitetyt tulokset osoittavat. Maatilan käyttöön tarkoitetuilla koneilla hoidettavien kenttäkokeiden suhteellisen suurilta ruuduilta otetut osanäytteet tulisi analysoida erikseen.