

## Results about $S_2$ -paracompactness

OHUD ALGHAMDI <sup>a</sup> AND LUTFI KALANTAN <sup>b</sup>

<sup>a</sup> Department of Mathematics, Faculty of Science and Arts in Almandaq, Al-Baha University, P.O.Box 1988, Al-Baha 65581, Saudi Arabia (ofalghamdi@bu.edu.sa)

<sup>b</sup> Department of Mathematics, King Abdulaziz University, Saudi Arabia (lkalantan@kau.edu.sa and lkalantan@hotmail.com)

Communicated by M. A. Sánchez-Granero

### ABSTRACT

---

We present new results regarding  $S_2$ -paracompactness, that we established in [1], and its relation with other properties such as  $S$ -normality, epinormality and  $L$ -paracompactness.

---

2020 MSC: 54C10; 54D20.

KEYWORDS: separable; paracompact;  $S$ -paracompact;  $S_2$ -paracompact;  $L$ -paracompact;  $L_2$ -paracompact;  $S$ -normal;  $L$ -normal.

### 1. INTRODUCTION

In this paper, we present some new results about  $S$ -paracompactness and  $S_2$ -paracompactness. First, we introduce the significant notations. An order pair will be denoted by  $\langle x, y \rangle$ . The sets of positive integers, rational numbers, irrational numbers and real numbers will be denoted by  $\mathbb{N}$ ,  $\mathbb{Q}$ ,  $\mathbb{P}$  and  $\mathbb{R}$ , respectively. The closure and the interior of the subset  $A$  of a topological space  $X$  will be denoted respectively by  $\overline{A}$  and  $\text{int}(A)$ . Throughout this paper, a  $T_1$  normal space is called  $T_4$  and a  $T_1$  completely regular space is called Tychonoff space ( $T_{3\frac{1}{2}}$ ). In the definitions of compactness, countable compactness, paracompactness, and local compactness we do not assume  $T_2$ . Moreover, in the definition of Lindelöfness we do not assume regularity. Also, the ordinal  $\gamma$  is the set of all ordinal  $\alpha$  such that  $\alpha < \gamma$ . We denote the first infinite ordinal by  $\omega$  and the first uncountable ordinal by  $\omega_1$ .

**Definition 1.1.** A topological space  $X$  is called  $S$ -paracompact if there exist a paracompact space  $Y$  and a bijective function  $f : X \rightarrow Y$  such that for every separable subspace  $A \subseteq X$  we have that  $f|_A : A \rightarrow f(A)$  is a homeomorphism. Moreover, if  $Y$  is  $T_2$  paracompact, then  $X$  is  $S_2$ -paracompact [1].

## 2. $S_2$ -PARACOMPACTNESS AND OTHER TOPOLOGICAL PROPERTIES

### 2.1. $S_2$ -paracompactness and $L_2$ -paracompactness.

Recall from [5] that a topological space  $X$  is called  $L$ -paracompact if there exist a paracompact space  $Y$  and a bijective function  $f : X \rightarrow Y$  such that  $f|_B : B \rightarrow f(B)$  is a homeomorphism for all Lindelöf subspace  $B$  of  $X$ . In addition, if  $Y$  is  $T_2$  paracompact, then  $X$  is  $L_2$ -paracompact.

Recall from [6] that a topological space  $X$  is called  $P$ -space if it is  $T_1$  and every  $G_\delta$  set is open. The countable complement topology defined on  $\mathbb{R}$ ,  $(\mathbb{R}, \mathcal{CC})$  (see [9, Example 20]), is an example of a space that is  $S_2$ -paracompact but not  $L_2$ -paracompact. It is  $S_2$ -paracompact because it is  $P$ -space, (see [1]), but not  $L_2$ -paracompact because it is Lindelöf and not paracompact space. In fact, it is not even  $L$ -paracompact.

We still do not have an answer for the following question:

Does there exist an  $L$ -paracompact space which is not  $S$ -paracompact?

**Theorem 2.1.** *If  $X$  is  $L$ -paracompact (resp.  $L_2$ -paracompact) such that for any separable subspace  $A \subseteq X$  there exists a Lindelöf subspace  $B \subseteq X$  such that  $A \subseteq B$ , then  $X$  is  $S$ -paracompact (resp.  $S_2$ -paracompact).*

*Proof.* Let  $X$  be  $L$ -paracompact such that for any separable subspace  $A \subseteq X$  there exists a Lindelöf subspace  $B \subseteq X$  such that  $A \subseteq B$ . Then, there exist a paracompact space  $Y$  and a bijective function  $f : X \rightarrow Y$  such that  $f|_B : B \rightarrow f(B)$  is a homeomorphism for every Lindelöf subspace  $B \subseteq X$ . Let  $A$  be any separable subspace of  $X$ . Then, there exists a Lindelöf subspace  $B$  of  $X$  such that  $A \subseteq B$ . Then,  $f|_A : A \rightarrow f(A)$  is a homeomorphism.  $\square$

A similar proof as in Theorem 2.1 yields the following corollaries.

**Corollary 2.2.** *If  $X$  is  $S$ -paracompact (resp.  $S_2$ -paracompact) such that for any Lindelöf subspace  $B \subseteq X$ , there exists a separable subspace  $A$  with  $B \subseteq A$ . Then,  $X$  is  $L$ -paracompact (resp.  $L_2$ -paracompact).*

Recall from [7] that a space  $X$  is called  $C$ -paracompact if there exist a paracompact space  $Y$  and a bijection  $f : X \rightarrow Y$  such that  $f|_K : K \rightarrow f(K)$  is a homeomorphism for every compact subspace  $K \subseteq X$ . Moreover, if  $Y$  is  $T_2$  paracompact, we say that  $X$  is  $C_2$ -paracompact.

**Corollary 2.3.** *If  $X$  is  $S$ -paracompact (resp.  $S_2$ -paracompact) such that for any compact subspace  $B \subseteq X$ , there exists a separable subspace  $A$  of  $X$  with  $A \subseteq B$ . Then,  $X$  is  $C$ -paracompact (resp.  $C_2$ -paracompact).*

**Application of Corollary 2.3:**

Take  $(\mathbb{R}, \mathcal{CC})$ , the countable complement topology on  $\mathbb{R}$ . Since  $A \subset \mathbb{R}$  is compact if and only if  $A$  is finite, we can say that every compact subspace is contained in a separable subspace of  $\mathbb{R}$ . Thus,  $(\mathbb{R}, \mathcal{CC})$  is  $C_2$ -paracompact.

Recall from [3, 4.4.F] that a space  $X$  is *locally separable* if each element of  $X$  has a separable open neighborhood.

**Theorem 2.4.** *Every  $S$ -paracompact (resp.  $S_2$ -paracompact) hereditarily locally separable is  $L$ -paracompact (resp.  $L_2$ -paracompact).*

*Proof.* Let  $X$  be  $S$ -paracompact (resp.  $S_2$ -paracompact) and hereditarily locally separable and let  $B$  be any Lindelöf subspace of  $X$ . Then,  $B$  is a locally separable Lindelöf subspace of  $X$ . Pick  $U_x$  to be a separable open neighborhood of each  $x \in B$ . Then,  $\{U_x\}_{x \in B}$  is an open cover of  $B$ . Let  $\mathcal{U}$  be a countable open subcover of  $\{U_x\}_{x \in B}$  and let  $D_x$  be a countable dense subset of each  $U_x \in \mathcal{U}$ . Then,  $D = \bigcup D_x$  is a countable dense subset of  $B$ , implying that  $B$  is separable. Therefore, since every Lindelöf subspace of  $X$  is separable and  $X$  is  $S$ -paracompact (resp.  $S_2$ -paracompact), then  $X$  is  $L$ -paracompact (resp.  $L_2$ -paracompact).  $\square$

**Problem 2.5.** *Does there exist a topological space that is  $L$ -paracompact (resp.  $L_2$ -paracompact) but not locally separable or not  $S$ -paracompact (resp.  $S_2$ -paracompact)?*

Note that local separability is essential in Theorem 2.4. For example,  $(\mathbb{R}, \mathcal{CC})$  is  $S$ -paracompact not locally separable. Observe that  $(\mathbb{R}, \mathcal{CC})$  is not  $L$ -paracompact.

**Theorem 2.6.** *Let  $X$  be a topological space such that the only separable or Lindelöf subspaces are the countable ones. Then,  $X$  is  $S$ -paracompact (resp.  $S_2$ -paracompact) if and only if  $X$  is  $L$ -paracompact (resp.  $L_2$ -paracompact).*

*Proof.* Let  $X$  be any topological space such that the only separable or Lindelöf subspaces are the countable ones. Suppose that  $X$  is  $S$ -paracompact (resp.  $S_2$ -paracompact). If  $B$  is any Lindelöf subspace of  $X$ , then  $B$  is countable, implying that  $B$  is separable. Hence,  $X$  is  $L$ -paracompact (resp.  $L_2$ -paracompact). Conversely, suppose that  $X$  is  $L$ -paracompact (resp.  $L_2$ -paracompact) and  $A$  is any separable subspace of  $X$ . Then,  $A$  is countable, implying that  $A$  is Lindelöf. Hence,  $X$  is  $S$ -paracompact (resp.  $S_2$ -paracompact).  $\square$

**Application of Theorem 2.6:**

Consider  $\omega_1$  with its usual ordered topology. Let  $A$  be any uncountable subset of  $\omega_1$ . Then  $A$  is not bounded. Hence,  $\{[0, \alpha] : \alpha < \omega_1\}$  is an open cover of  $A$  that has no countable subcover, which implies that  $A$  is not Lindelöf. Since  $\omega_1$  satisfies the condition in Theorem 2.6, then  $\omega_1$  is  $L_2$ -paracompact because it is  $S_2$ -paracompact, (see [1]).

A family  $\{A_s\}_{s \in S}$  of subsets of a space  $X$  is called point-finite if for each  $x \in X$ , the set  $\{s \in S : x \in A_s\}$  is finite, (see [3]).

Recall from [9] that a space  $X$  is *metacompact* if every open cover of  $X$  has a point-finite open refinement.

**Theorem 2.7.** *Any hereditarily metacompact  $L$ -paracompact (resp.  $L_2$ -paracompact) is  $S$ -paracompact (resp.  $S_2$ -paracompact).*

*Proof.* Let  $X$  be  $L$ -paracompact (resp.  $L_2$ -paracompact) hereditarily metacompact and let  $A$  be any separable subspace of  $X$ . Then,  $A$  is a separable metacompact subspace of  $X$ . Suppose that  $A$  is not Lindelöf. Then, there exists an open cover of  $A$ , say  $\mathcal{W} = \{W_\alpha : \alpha \in \Lambda\}$ , which has no countable subcover. Let  $\mathcal{U}$  be a point-finite open refinement of  $\mathcal{W}$ . Then,  $\mathcal{U}$  is uncountable by our assumption. Let  $D$  be the countable dense subset of  $A$ . Hence,  $D \cap U \neq \emptyset$  for all  $U \in \mathcal{U}$  implying that there exists  $d \in D$  contained in uncountable members of  $\mathcal{U}$  which contradicts the fact that  $\mathcal{U}$  is a point-finite family. Hence,  $A$  is Lindelöf, implying that  $X$  is  $S$ -paracompact (resp.  $S_2$ -paracompact).  $\square$

**Problem 2.8.** *Does there exist a topological space which is  $S$ -paracompact (resp.  $S_2$ -paracompact) but not hereditarily metacompact or  $L$ -paracompact (resp.  $L_2$ -paracompact)?*

2.2.  $S_2$ -paracompactness and Epinormality.

**Definition 2.9.** A topological space  $(X, \tau)$  is *epinormal* if there exists a coarser topology, say  $\mathcal{V}$ , such that  $(X, \mathcal{V})$  is  $T_4$ , (see [2]).

Since every epinormal space is Hausdorff as it is proved in [2], then the countable complement topology on  $\mathbb{R}$ ,  $(\mathbb{R}, \mathcal{CC})$ , is an example of  $S_2$ -paracompact that is not epinormal. On the other hand, the following example shows that there exists an epinormal space which is not  $S_2$ -paracompact.

**Example 2.10.** Let  $A = \{\langle x, 0 \rangle : 0 < x \leq 1\}$  and  $B = \{\langle x, 1 \rangle : 0 \leq x < 1\}$ .




FIGURE 1. This figure illustrates the neighborhood system of Strong Parallel Line Topology  $(X, \sigma)$ .

The *strong parallel line* topology  $\sigma$  on  $X = A \cup B$  is the unique topology generated by the following neighborhood system:

For each  $\langle t, 0 \rangle \in A$ , let  $\mathfrak{B}(\langle t, 0 \rangle) = \{U : U = \{\langle x, 0 \rangle : 0 \leq a < x \leq t\} \cup \{\langle x, 1 \rangle : a < x < t\}\}$ , and for each  $\langle t, 1 \rangle \in B$ , let  $\mathfrak{B}(\langle t, 1 \rangle) = \{V : V = \{\langle x, 1 \rangle : t \leq x < b \leq 1\}\}$ , (see [9, Example 96]).

Since  $(X, \sigma)$  is separable and not paracompact space because it is a Hausdorff and not regular topological space, then  $(X, \sigma)$  cannot be  $S_2$ -paracompact.

Define  $\tau$  on  $X$  to be the unique topology that is generated by the following neighborhood system:

Every element in  $A$  has the same local base as  $\sigma$  and for each element  $\langle t, 1 \rangle \in B$ , let  $\mathfrak{B}(\langle t, 1 \rangle) = \{V : V = \{\langle x, 0 \rangle : t < x < b \leq 1\} \cup \{\langle x, 1 \rangle : t \leq x < b\}\}$ . The topology  $(X, \tau)$  is named *weak parallel line*, (see [9, Example 96]).




FIGURE 2. This figure illustrates the neighborhood system of Weak Parallel Line Topology  $(X, \tau)$ .

Define a relation  $\preceq$  on  $X$  as follows:

For  $\langle x, y \rangle$  and  $\langle k, l \rangle \in X$ , we write  $\langle x, y \rangle \preceq \langle k, l \rangle$  if and only if either  $x < k$  or  $x = k$  and  $y = 0 < l = 1$ , or  $x = k$  and  $y = l$ . Then,  $(X, \tau)$  is a linearly ordered topological space (LOTS). Since any LOTS is  $T_4$ , we have  $(X, \tau)$  is  $T_4$  and  $\tau$  is coarser than  $\sigma$ , hence, we get that  $(X, \sigma)$  is epinormal.

**Theorem 2.11.** *Any  $S_2$ -paracompact Fréchet space is epinormal.*

*Proof.* Let  $(X, \tau)$  be  $S_2$ -paracompact Fréchet space. Without loss of generality, assume that  $(X, \tau)$  is not normal. Let  $(Y, \tau')$  be a  $T_2$  paracompact space and let  $f : X \rightarrow Y$  be a bijective function such that  $f|_A : A \rightarrow f(A)$  is a homeomorphism for every separable subspace  $A \subseteq X$ . Then,  $f$  is continuous since  $X$  is Fréchet. Consider  $\mathcal{V} = \{f^{-1}(U) : U \in \tau'\}$ . Then,  $\mathcal{V}$  is a topology on  $X$  and since any open set in  $\mathcal{V}$  is open in  $\tau$  by continuity of  $f$ , we get that  $\mathcal{V}$  is coarser than  $\tau$ . Observe that  $f : (X, \mathcal{V}) \rightarrow (Y, \tau')$  is a homeomorphism. Therefore,  $(X, \mathcal{V})$  is  $T_2$  paracompact and, hence,  $T_4$ .  $\square$

For the converse of Theorem 2.11, we have the left ray topological space defined on  $\mathbb{R}$ ,  $(\mathbb{R}, \mathcal{L})$ , as an example of epinormal Fréchet space that is not  $S_2$ -paracompact since it is separable and not paracompact space.

### 2.3. $S_2$ -paracompactness and $S$ -normality.

Recall that a topological space  $X$  is  $S$ -normal if there exist a normal space  $Y$  and a bijective function  $f : X \rightarrow Y$  such that  $f|_A : A \rightarrow f(A)$  is a homeomorphism for each separable subspace  $A$  of  $X$ , (see [4]). From the definition of  $S$ -normality, it is clear that any  $S_2$ -paracompact is  $S$ -normal. However, we show in the following example that this relation is not reversible.

**Example 2.12.** An example of a  $T_4$  topological space that is  $S$ -normal but not  $S_2$ -paracompact is the sigma product  $\Sigma(0)$  as a subspace of  $2^{\omega_1}$ , where  $2 = \{0, 1\}$  considered with the discrete topology. It is not  $S_2$ -paracompact since it cannot be condensed onto a  $T_2$  paracompact space, (see [8]).

**Theorem 2.13.** *Let  $X$  be Fréchet and Lindelöf space such that any finite subspace of  $X$  is discrete.  $X$  is  $S$ -normal if and only if  $X$  is  $S_2$ -paracompact.*

*Proof.* Let  $Y$  be a normal space and let  $f : X \rightarrow Y$  be a bijective function such that  $f|_A : A \rightarrow f(A)$  is a homeomorphism for each separable subspace  $A$  of  $X$ . Without loss of generality, let  $X$  have more than one element. Thus,  $Y$  is  $T_1$  since any finite subspace of  $X$  is separable and discrete. By continuity of  $f$  and since  $X$  is Lindelöf, then  $Y$  is Lindelöf. Since  $Y$  is  $T_3$  Lindelöf, then  $Y$  is  $T_2$  paracompact. Thus,  $X$  is  $S_2$ -paracompact.

Conversely, assume that  $X$  is  $S_2$  paracompact. Let  $Y$  be a  $T_2$  paracompact space and let  $f : X \rightarrow Y$  be a bijective function such that  $f|_A : A \rightarrow f(A)$  is a homeomorphism for each separable subspace  $A$  of  $X$ . Hence, since  $Y$  is  $T_2$  paracompact, then  $Y$  is  $T_4$ . Therefore,  $X$  is  $S$ -normal.  $\square$

Recall that from [3] that a topological space  $X$  is *locally metrizable* if there exists a metrizable open neighborhood for each  $x \in X$ .

**Theorem 2.14.** *If  $X$  is hereditary Lindelöf,  $S_2$ -paracompact and locally metrizable, then  $X$  is  $T_2$  paracompact and, hence,  $T_4$ .*

*Proof.* Set  $U_x$  to be a metrizable open neighborhood of each  $x \in X$ . Since  $X$  is Lindelöf, then there exists a countable set  $E$  such that  $X \subseteq \bigcap_{x \in E} U_x$ . Now, since  $X$  is hereditary Lindelöf, then  $U_x$  is Lindelöf as a subspace of  $X$  for every  $x \in X$ . Hence,  $U_x$  is separable being Lindelöf and metrizable for every  $x \in X$ . Since  $X$  is  $S_2$ -paracompact and separable, then  $X$  is  $T_2$  paracompact and, hence,  $T_4$ .  $\square$

Let  $(X, \tau)$  be a topological space and let  $M$  be a proper nonempty subset of  $X$ . The discrete extension of the topological space  $(X, \tau)$  is defined by the following neighborhood system:

For each  $x \in X \setminus M$ , let  $\mathcal{B}(x) = \{\{x\}\}$  and for each  $x \in M$ , let  $\mathcal{B}(x) = \{U \in \tau : x \in U\}$ . We denote the discrete extension of  $X$  by  $X_M$ , (see [9]).

The following example shows that the discrete extension of  $S_2$ -paracompact need not to be  $S_2$ -paracompact.

**Example 2.15.** Consider  $(\mathbb{R}, \mathcal{RS})$ , the rational sequence topology on  $\mathbb{R}$ , (see [9, Example 65]). Since  $\mathcal{RS}$  is separable and not paracompact, then it is not  $S_2$ -paracompact. Also, because it is a Tychonoff locally compact space, we can set  $X = \mathbb{R} \cup \{p\}$  to be the one point compactification of it.  $X$  is  $T_2$  compact, which implies that  $X$  is  $S_2$ -paracompact. Consider  $X_{\mathbb{R}}$ , the discrete extension of  $X$ . Since  $\{p\}$  is closed and open subset in  $X_{\mathbb{R}}$ , then  $\mathbb{R}$  is a closed subspace of  $X_{\mathbb{R}}$ . However, since  $(\mathbb{R}, \mathcal{RS})$  is not normal, we conclude that  $X_{\mathbb{R}}$  cannot be normal. Since  $X_{\mathbb{R}}$  is  $T_2$  and not normal space, then  $X_{\mathbb{R}}$  is not paracompact. Since  $X_{\mathbb{R}}$  is separable as  $\mathbb{Q} \cup \{p\}$  is a countable dense subset of  $X_{\mathbb{R}}$ , then  $X_{\mathbb{R}}$  is not  $S_2$ -paracompact.

## REFERENCES

- [1] O. Alghamdi, L. Kalantan and W. Alagal,  $S$ -paracompactness and  $S_2$ -paracompactness, *Filomat* 33, no. 17 (2019), 5645–5650.
- [2] S. Alzharani and L. Kalantan, Epinormality, *J. Nonlinear Sci. Appl.* 9 (2016), 5398–5402.
- [3] R. Engelking, *General Topology*, PWN, Warszawa, 1977.
- [4] L. Kalantan and M. Alhomieyed,  $S$ -normality, *J. Math. Anal.* 9, no. 5 (2018), 48–54.
- [5] L. Kalantan,  $L$ -paracompactness and  $L_2$ -paracompactness, *Hacet. J. Math. Stat.* 48, no. 3 (2019), 779–784.
- [6] A. K. Misra, A topological view of  $P$ -spaces, *Gen. Topol. Appl.* 2 (1972), 349–362.
- [7] M. M. Saeed, L. Kalantan and H. Alzumi,  $C$ -paracompactness and  $C_2$ -paracompactness, *Turk. J. Math.* 43 (2019), 9–20.
- [8] M. M. Saeed, L. Kalantan and H. Alzumi, Result about  $C_2$ -paracompactness, *Eur. J. Pure Appl. Math.* 14, no. 2 (2021), 351–357.
- [9] L. Steen and J. A. Seebach, *Counterexamples in Topology*, Dover Publications, 1995.
- [10] V. Zaitsev, On certain classes of topological spaces and their bicomactification, *Dokl. Akad. Nauk SSSR* 178 (1968), 778–779.