

SEVERITY EVALUATION METHODS FOR SOUTHERN POLYSORA CORN RUST

MÉTODOS DE AVALIAÇÃO DA SEVERIDADE DA FERRUGEM POLYSORA DO MILHO

Fernando Cezar JULIATTI¹; Adriana Andrade FIGUEIRÓ²; Élcio Oliveira ALVES³

1. Professor, Doutor, Instituto de Ciências Agrárias – ICIAG, Universidade Federal de Uberlândia, Uberlândia, MG, Brasil; 2. Bióloga, Mestre em Fitopatologia e Doutora em Fitotecnia pela UFRGS. Pós Doutorado-CAPEL, LAMIP-ICIAG-UFU, Uberlândia, MG, Brasil; 3. Engenheiro Agrônomo, Pioneer Sementes, Itumbiara, GO, Brasil

ABSTRACT: Among the biotrophic organisms that attack the corn crop the most important for Central Brazil is the *Puccinia polysora* Underw, which is the causal agent of polysora corn rust. Due to the need to study the methods used in the assessments better, this work was performed. An experiment to identify the best methodology for assessing polysora rust was set up in Itumbiara – Goiás - Brazil, with 22 hybrids and an experimental randomized block design with three replications. So as to evaluate disease severity in relation to the leaf area affected, the Area Under the Disease Progress Curve (AUDPC) and a single evaluation were used, while also considering the severity of the disease on the leaf area affected, 30 days after flowering (30 d.a.f.) Both based on a proposal diagrammatic scale from Agrocerees with grades ranging from one (highly resistant) to nine (highly susceptible), which were employed considering the plot as a whole. So as to calculate the AUDPC five evaluations of disease severity at intervals of 10 days which begun 50 days after sowing were carried out. It was found that it was possible to determine the level of genotype resistance by using the AUDPC or by using the single assessment at 30 d.a.f., with the assessment of disease severity by means of the AUDPC calculation allowing for better understanding of the evolution of the disease in time and that the one only or single assessment at 30 d.a.f. allows us to work with a greater number of genotypes, due to the practicality of the methodology.

KEYWORDS: *Puccinia polysora*. Evaluation. Disease. Rust.

INTRODUCTION

Since the beginning of the 1980s, polysora rust, *Puccinia polysora* Underw, has been a serious problem in the corn crop in south-west regions of Goiás, in the Triângulo Mineiro and more recently in the northwest of São Paulo, in the East and in the North of Paraná and in Mato Grosso do Sul in Brazil. It is considered to be the most aggressive and destructive among corn diseases and can cause rapid necrosis of the plant. Under favorable conditions and in susceptible cultivars, this rust may occur with severe presence in the leaves, stems, straw, and hence cause a marked reduction in the size of the corn cobs and grains (FANTIN, 1997).

The severity of the disease can be assessed by subjective methods such as counting the number of lesions per leaf or the measurement of the affected area, but these methods are usually very time consuming. The quantification of disease severity can be carried out visually by means of numeric scales, grades or percentage and diagrammatic procedures (JULIATTI; SANTOS, 1999).

As corn breeding or improvement programs are very dynamic, and produce a large number of new varieties each year, it is necessary to evaluate the behavior of these materials in relation to this

disease. This, not only so as to direct future work of breeding/improvement, in order to obtain resistant cultivars, but also to act as a guide in the selection and recommendation of cultivars for different regions (VON PINHO et al., 2001).

The less costly and more efficient control method for polysora rust is the use of hybrids or varieties with satisfactory levels of resistance to the pathogen (BALMER; PEREIRA, 1987; McGEE, 1988; PEREIRA, 1997; PINTO et al., 1997; SHERF; MACNAB, 1986). The objective of this work was to compare the effectiveness of methods used in the severity analysis of polysora corn rust.

MATERIAL AND METHODS

The trial was conducted at the Pioneer Seeds Ltd. Research Center, in Itumbiara GO, in the period from October 29, 2003 to March 10, 2004. The altitude at the location is 511 m, at coordinates S18° 19.995 and W49° 12.101 and the soil is of medium texture (sandy clay), with an approximate slope of 1%. The average annual rainfall is 1,450 mm, with a minimum annual average temperature of 20 °C and maximum 30 °C.

The preparation of the area was carried out using a disk harrow (26" disks) and then a leveling harrow to complete the preparation.

The fertilization of the planting area was made with 500 kg.h⁻¹ of 08-18-20 formula +Zn, based on previous analysis of the soil. The topdressing was carried out using 120 kg of N divided into two applications of urea (44% N), at 25 and 35 days after sowing (d.a.s.), respectively.

The pre-emergent herbicide Agimix (Atrazine 26% + Metolachlor 26%) was used at a dose rate of 5 L.ha⁻¹ for weed control. For insect control the Lorsban BR480 products (Chlorpirifos 48%) were used at the dosages specified by the manufacturer, at V3 and V5 phenological stages, and Lannate BR (Methomil 21.5%) product, also at the dosage recommended by the manufacturer at V8 and V10 phenological stages.

The experimental design was that of randomized blocks, combining 22 hybrids with different levels of genetic resistance, randomly distributed in three blocks (Table 1).

Seeds were sown on October 29th 2003. The plots were formed by three lines of 5 m, spaced at 0.75 m between rows, with a total used area of 11.25 m per plot and 495 m² of total experiment area. To evaluate the severity of the polysora rust and harvesting the overall plot was considered.

The *P. polysora* inoculum was obtained from urediniospores collected from the 30F53 hybrid, a hybrid not adapted to the region and highly susceptible to the pathogen. The urediniospores were collected using a paintbrush and a tray which was lined with foil and were subsequently packed in gelatin capsules until the time of inoculation, carried out on the same day. The inoculum was prepared at a concentration of 1x10⁵ urediniosporos.mL⁻¹ solution plus 0,05 µL.mL⁻¹ of Tween 20; thereupon the inoculum was homogenized for 10 minutes.

Table 1. Agronomic characteristics of the hybrids used in the test.

Identification of Hybrid	Type of Hybrid	Cycle	Company	Grain Type
AG7000	Simple	Average	Agroceres	Medium hard
AG7575	Simple	Early	Agroceres	Medium hard
2C577	Simple	Early	Dow AgroSciences	Semi-dent
2C599	Simple	Early	Dow AgroSciences	Medium hard
DKB350	Triple	Average	Monsanto	Medium hard
30F53	Simple	Early	Pioneer	Medium hard
30P70	Simple	Early	Pioneer	Medium hard
30R50	Simple	Early	Pioneer	Medium hard
DKB390	Simple	Average	Monsanto	Medium hard
30F80	Simple	Semi-early	Pioneer	Hard
30F88	Simple	Semi-early	Pioneer	Hard
30F90	Simple	Semi-early	Pioneer	Hard
30K75	Simple	Semi-early	Pioneer	Medium hard
Fort	Simple	Early	Syngenta	Hard
Strike	Simple	Early	Syngenta	Hard
Tractor	Double	Early	Syngenta	Hard
Valent	Triple	Early	Syngenta	Hard
AG9010	Simple	Super-early	Monsanto	Hard
30F33	Simple	Early	Pioneer	Hard
30F44	Simple	Early	Pioneer	Medium hard
Speed	Simple	Super-early	Syngenta	Hard
3081	Simple	Super-early	Pioneer	Hard

The inoculation was carried out in borders containing the highly susceptible hybrid, 30F53, at the V8-V10 stage, about 20 days before flowering, so that these function as disseminator lines of the pathogen.

To estimate the Area Under Disease Progress Curve (AUDPC) five evaluations of disease severity were performed. The first evaluation was performed 50

days after sowing (d.a.s.), the others at 60, 70, 80 and 90 d.a.s., respectively.

For the evaluations the total experimental plot system was used, which consisted in assessing disease severity in relation to the percentage of leaf area affected of all the experimental plot, according to the diagrammatic scale of Agroceres Health Guide (AGROCERES, 1996) (Table 2).

Table 2. Leaf area affected by the disease, and the corresponding score according to the AGROCERES (1996) diagrammatic scale.

Original Score	Leaf Area (%)
9	100
8	90
7	80
6	70
5	50
4	30
3	10
2	0,5
1	0

The AUDPC was calculated from the disease progress curve, based on the percentage of leaf area affected in the experimental plot:

$$AUDPC = \sum_{i=1}^{n-1} \frac{(Y_{i+1} + Y_i)}{2} \times (T_{i+1} - T_i)$$

In which: Y_i : disease severity at the time of evaluation i ($i = 1, \dots, n$), Y_{i+1} : disease severity at time of evaluation $i + 1$, T_i : time of evaluation i (number of days after plant emergence), T_{i+1} : time of evaluation $i + 1$ and n : total number of observations.

For AUDPC data obtained in the plot, the statistical model used for the variance analysis was as follows, considering all effects as fixed, except the experimental error:

$$Y_{ij} = m + g_i + b_j + e_{ij}$$

In which: Y_{ij} : AUDPC of genetic material i in block j , m : average overall effect, g_i : effect of genetic material ($i = 1, 2, \dots, 22$), b_j : effect of block j ($j = 1, \dots, 3$) and e_{ij} : effect of experimental error.

To determine the severity of the disease in a single assessment the assessment at 30 days after the average flowering (d.a.f.) of the plot was considered.

Harvesting was carried out using a Wintersteiger harvester in which data as to grain

weight and moisture were collected simultaneously. The whole plot was considered for harvesting. Data were analyzed using the SANEST¹ program for the F test, and average (Tukey) test.

For the yield data obtained at plot level, the statistical model used for variance analysis was as follows, considering all effects as fixed, except the experimental error:

$$Y_{ij} = m + g_i + b_j + e_{ij}$$

In which: Y_{ij} : Productivity of genetic material i in block j , m : average overall effect, g_i : effect of genetic material ($i = 1, 2, \dots, 22$), b_j : effect of block j ($j = 1, \dots, 3$) and e_{ij} : effect of experimental error.

Resistance classification by AUDPC was based on Table 3, which was established after analysis of data obtained in experiment.

Resistance classification by severity at 30 days after flowering (d.a.f.) was based on Table 6.

The average maximum temperature in the culture over the period was approximately 30°C and the average minimum was about 20 °C. The rainfall index was high, which is characteristic of the region driving the period when the test was being carried out (Figure 1).

Table 3. Resistance classification by AUDPC (Area Under Disease Progress Curve).

AUDPC	Resistance Classification*
1 a 20	HS
21 a 200	R
201 a 500	MR
501 a 1000	MS
1001 a 2000	S
>2000	HS

HS: highly susceptible, S: susceptible, MS: moderately susceptible, MR: moderately resistant, R: resistant and HR: highly resistant.




Figure 1. Climatological data from the Research Center of Pioneer Seeds Ltd, from October 2003 to March 2004.

RESULTS AND DISCUSSION

At 50 d.a.s. only some/slight symptom of polysora rust was noted in the trial. The hybrids which at this stage showed symptoms were later the most susceptible and were classified as highly susceptible both by the AUDPC calculation as well as by assessing the severity of the disease in experimental plot at 30 d.a.f. (Table 4).

There was a great uniformity in the severity of polysora rust among all plants of the plot, allowing for an optimal evaluation of the disease both by AUDPC as well as by assessing the severity at a single time, at 30 d.a.f.. Studies indicate that both methods were also effective in discriminating the level of resistance to *P. polysora* in corn hybrids, allowing for classification in a similar manner (Von Pinho et al., 2000).

The 22 hybrids used in this trial showed different levels of resistance to the pathogen (Table 1). The genotypes that presented themselves as highly susceptible to disease are not indicated by the companies responsible for planting in the environment where the trials were installed, these genotypes just were part of the experiment for the study.

Both methods used in the evaluation enabled the classification of the materials in a

similar manner (Table 4). However, the once only or single evaluation at 30 d.a.f. proved to be efficient and more practical, due to the possibility of evaluating a larger number of genotypes in a shorter time frame.

The highly susceptible hybrids and those susceptible to disease presented the first symptoms at 50 d.a.s. (Figure 2), with some uredia on the lower leaves. For the highly susceptible genotypes, the symptoms showed an upward development after 50 d.a.s., colonizing the whole host by the 90 d.a.s. The susceptible hybrids showed an upward development of symptoms from the 70 d.a.s. colonizing an average of 70% of the host by the 90 d.a.s. (Figure 2).

The moderately resistant genotypes presented symptoms on lower leaves a little before 60 d.a.s. No upward development of the disease as in highly susceptible genotypes was verified, but the symptoms intensified after 70 d.a.s., as in the susceptible genotypes (Figure 2).

The resistant or highly resistant genotypes to the pathogen presented few leaf symptoms, only after 70 d.a.s. and did not develop these symptoms in the upper leaves. Symptoms were limited to the first three lower leaves, without causing visually observed damage (Figure 2).




Figure 2. Temporal evolution of polysora rust in different resistance levels presented by the genetic material HS (highly susceptible), S (susceptible), MS (moderately susceptible), MR (moderately resistant), R (resistant) and HR (highly resistant) from the calculation of AUDPC. AUDPC evaluated at 40, 50, 60, 70, 80 and 90 days after sowing. Ver cores das linhas em caso de publicação em tons de cinza

Some genotypes showed small differences in classification between the two evaluation methods (Table 4). The AG9010 genotype was classified as resistant by AUDPC and moderately

resistant by the evaluation at 30 d.a.f. In this genotype there was an increase in disease severity between 80 and 90 d.a.f. (Figure 3).




Figure 3. Comparison between AUDPC (Area Under Disease Progress Curve) of AG9010 with other hybrids with different levels of resistance. AUDPC evaluated at 40, 50, 60, 70, 80 and 90 days after sowing. Ver cores das linhas em caso de publicação em tons de cinza

When comparing AG9010 with other genotypes with different resistance levels (3081 – AS, 30F80 – MR, AG7000 – R, DKB350, R e 30F90 – AR) it was found that AG9010 had a behavior similar to the highly susceptible genotypes from the 80 d.a.s. (Figure 2 and 3).

The Tractor and 30K75 genotypes obtained different resistance classification by both severity assessment methods (Table 4). The Tractor hybrid

was classified as susceptible by AUDPC and as moderately susceptible by severity at 30 d.a.f. The 30K75 hybrid was classified as moderately resistant by AUDPC and as moderately susceptible by severity. It can be observed by AUDPC values and severity at 30 d.a.f. that both are located in an area of transition between the resistance groups that were classified, as well as AG9010 (Table 4). AG9010, which was classified as resistant by AUDPC and

moderately resistant by the evaluation at 30 d.a.f., produced 5.273 kg.ha⁻¹ (Table 5). This showed that even with a good level of resistance, this genotype showed a limited production capacity.

The 30P70, 30F44 and Tractor genotypes were classified as susceptible to *P. polysora* for symptoms of the disease and produced 6,590, 5,650 and 5.678 kg.ha⁻¹, respectively (Table 5). Therefore,

while even presenting high levels of disease severity, these genotypes were highly productive.

The 30F90 genotype was highly resistant to fungus and had high productivity, presenting a high adaptability to the environment in lowland Central Brazil with or without the presence of disease (Tables 4 and 5).

Table 4. Severity of *P. polysora* on the AUDPC (Area Under Disease Progress Curve), resistance and severity classification in the experimental plot at 30 d.a.f. (days after flowering).

Genotypes	AUDPC ¹	Resistance AUDCP ²	Sev.30 d.a.f. ³	Resistance Sev.30 d.a.f. ⁴
30F53	2807.0 a*	HS	100.0 a*	HS
30R50	2807.0 a	HS	100.0 a	HS
3081	2054.7 b	HS	100.0 a	HS
30F44	1238.0 c	S	76.7 ab	S
30P70	1188.0 c	S	76.7 ab	S
Tractor	1043.0 cd	S	56.7 bcd	MS
DKB390	707.3 cde	MS	30.0 def	MS
Speed	679.0 cde	MS	63.3 bc	MS
30F33	578.7 cde	MS	23.3 efg	MR
Valent	429.0 de	MR	23.3 efg	MR
30K75	415.0 de	MR	36.7 cde	MS
Fort	393.0 de	MR	16.7 efg	MR
2C577	348.3 de	MR	16.7 efg	MR
30F80	312.3 de	MR	23.3 efg	MR
30F88	149.0 e	R	6.7 fg	R
AG7000	144.0 e	R	6.7 fg	R
AG9010	112.3 e	R	10.7 efg	MR
DKB350	80.7 e	R	3.3 fg	R
AG7575	73.3 e	R	3.3 fg	R
30F90	17.3 e	HR	0.0 g	HR
2C599	14.7 e	HR	0.0 g	HR
Strike	12.3 e	HR	0.0 g	HR

*Averages followed by different letters differ from one another at the level of significance of 1% of probability by the Tukey test; AUDPC: d.m.s. 5% = 637.70835, d.m.s. 1% = 735.99692; Severity: d.m.s. 5% = 23.85753, d.m.s. 1% = 27.53464; ¹Averages of AUDPC (Area Under the Disease Progress Curve); ²Resistance classification according to AUDPC data, with: HS: highly susceptible, S: susceptible, MS: moderately susceptible, MR: moderately resistant, R: resistant and HR: highly resistant; ³ Averages of disease severity assessed at 30 daf. ⁴ Resistance classification according to the severity of disease data assessed at 30 d.a.f. (days after flowering).

Table 5. Resistance classification (AUDPC and evaluation at 30 d.a.f.) and productivity (kg.ha⁻¹) of corn genotypes.

Genotypes	Resistance 30 d.a.f.	Resistance AUDPC ²	Productivity (kg.ha ⁻¹) ³	
30F53	HS	HS	4301.6 cd**	B*
30R50	HS	HS	4481.3 bcd	B
3081	HS	HS	4020.6 d	B
30F44	S	S	5652.6 abcd	AB
30P70	S	S	6592.3 abcd	AB
TRACTOR	MS	S	5678.0 abcd	AB
DKB390	MS	MS	6398.0 abcd	AB
SPEED	MS	MS	6153.6 abcd	AB

30F33	MR	MS	6345.6 abcd	AB
VALENT	MR	MR	5700.0 abcd	AB
30K75	MS	MR	6493.0 abcd	AB
FORT	MR	MR	6348.3 abcd	.AB
2C577	MR	MR	6371.3 abcd	.AB
30F80	MR	MR	5749.3 abcd	AB
30F88	R	R	6018.6 abcd	AB
AG7000	R	R	6310.0 abcd	AB
AG9010	MR	R	5273.3 abcd	AB
DKB350	R	R	6398.3 abcd	AB
AG7575	R	R	6097.3 abcd	AB
30F90	HR	HR	7805.3 a	A
2C599	HR	HR	6860.3 abc	AB
STRIKE	HR	HR	7003.0 ab	AB

Average 5667,3

Averages followed by different letters differ from one another at the significance level of 1%* or 5%** of probability by the Tukey test; AUDPC: d.m.s. 5% = 637.70835, d.m.s. 1% = 735.99692; Productivity: d.m.s. 5% = 2651.33333, d.m.s., 1% = 3059.9774; Variation Coefficient of Productivity: 14%; ²Resistance classification according to AUDPC data (HS: highly susceptible, S: susceptible, MS: moderately susceptible, MR: moderately resistant, R: resistant and HR: highly resistant); ³Productivity (kg.ha⁻¹).

Using the correlation of productivity with the severity of the disease evaluated by AUDPC it was possible to observe that there was a negative relationship between productivity and AUDPC of 60% ($r = -0,60$) (data not shown).

The AUDPC correlation with the assessment of severity in a single assessment at 30 d.a.f., was 87% (data not shown). It is considered, therefore, that as much by means of the AUDPC as by means of the single evaluation after 30 d.a.f., the selection of genotype for resistance to *P. polysora* is possible. Therefore, the evaluation of severity at 30 d.a.f. enables the discrimination of genotypes, reducing labor and time available for this operation, as regards the evaluation by AUDPC. These results corroborate the data found by Von Pinho et al. (2000).

Foss (1999), working with the assessment of damage caused by *P. polysora* in corn, found that the experimental plot method with stand correction was satisfactory for the evaluation of the damage caused by polysora rust in corn production. There is a good correlation between the severity of the disease and AUDPC with production. In relation the popcorn diseases significant differences ($P \leq 0.01$) of resistance to southern rust (*Puccinia*

polysora) northern leaf blight (*Exserohilum turcicum*) and Phaeosphaeria leaf spot (complex *Phaeosphaeria maydis/ Pantoaea ananatis*) were observed. The cluster analysis detected different resistance levels to the series of leaf diseases. In 12 of the new hybrids the resistance level to the set of diseases was similar as in the commercial hybrid IAC 112, considered the best reference for leaf disease resistance among commercial popcorn genotypes (VIEIRA et al. 2009).

CONCLUSIONS

The assessment of severity by way of AUDPC and by evaluation at 30 d.a.f. allows one to characterize and classify corn genotypes for resistance to *P. polysora*.

The assessment by AUDPC enables the visualization of the progression of the disease in time for the different levels of resistance of the genotypes. However, the single assessment at 30 d.a.f. was more practical and feasible when one has a large number of genotypes to analyze.

RESUMO: Dentre os organismos biotróficos que atacam a cultura do milho o mais importante para o Brasil Central é a *Puccinia polysora* Underw, agente causal da ferrugem polysora do milho. Devido à necessidade de se estudar melhor os métodos utilizados nas avaliações, realizou-se este trabalho. Um experimento para identificar a melhor metodologia para avaliação da ferrugem polysora foi montado em Itumbiara – Goiás, com 22 híbridos e o delineamento experimental de blocos casualizados com três repetições. Utilizou-se para as avaliações da severidade da doença em relação à área foliar afetada, a Área Abaixo da Curva de Progresso da Doença (AACPD) e a avaliação única, considerando também a severidade da doença em relação à área foliar afetada, aos 30 dias após florescimento (30 d.a.f.). Ambas

baseadas na escala diagramática da Agroceres, com notas variando de um (altamente resistente) a nove (altamente suscetível), que foi empregada considerando a parcela com um todo. Para o cálculo da AACPD foram realizadas cinco avaliações da severidade da doença a intervalos de 10 dias e iniciadas aos 50 dias após semeadura. Verificou-se que tanto pela AACPD ou pela avaliação única aos 30 d.a.f. foi possível a determinação do nível de resistência dos genótipos, sendo que a avaliação da severidade da doença através do cálculo da AACPD permite entender melhor a evolução da doença no tempo e a avaliação única aos 30 d.a.f. nos permite trabalhar com um maior número de genótipos, devido à praticidade da metodologia.

PALAVRAS-CHAVE: *Puccinia polysora*. Avaliação. Doença. Ferrugem.

REFERENCES

- AGRIOS, G. N. **Plant pathology**. 4. ed. New York: Academic. New York, 1998. 635p.
- AGROCERES. **Guia Agroceres de sanidade**. São Paulo: 1996. 72p.
- BOLOW, J. F. W. As ferrugens (*P. sorghi*, *P. polysora*, *Physopella zae* do milho (*Zea mays*). I.Revisão bibliográfica. **Pesquisa Agropecuária Brasileira**, Brasília, DF, v. 1, p.249-262, 1966.
- BONDE, M. R.; BROMFIELD, K. R.; MELCHING, J. S. Morfological development of *Physopella zae* on corn. **Phytopathology**, Saint Paul, v. 72, n. 11, p.1489-1491, 1982. <http://dx.doi.org/10.1094/Phyto-72-1489>
- CAB INTERNATIONAL. **Crop Protection Compendium**. Global Module. 2003.ed. Wallingford, 2003.
- FANCELLI, A.L. **Ecofisiologia e Fenologia**. CAD – Tecnologia da Produção de Milho. Piracicaba: Aldeia Norte, São Paulo. 2002. 52p.
- FANTIN, G. M. **Avaliação de resistência do milho à ferrugem causada por *Puccinia polysora* Underw.** Piracicaba. 1997. 136p.
- FERNANDES, F.T.; OLIVEIRA, E.. **Principais doenças na cultura do milho**. Sete Lagoas: EMBRAPA-CNPMS, 2000. 80p.
- FOSS, M.R.D. **Avaliação de danos causados por *Puccinia polysora* em milho (*Zea mays*)**. 2000. 67f. Dissertação (Mestrado em Genética e Melhoramento de Plantas) – Faculdade de Ciências Agrárias e Veterinárias, Universidade Estadual Paulista “Júlio de Mesquita Filho”, Jaboticabal.
- GODOY, C. V. et al. Influência da temperatura e do período de molhamento nos parâmetros monocíclicos de *Puccinia polysora*. **Fitopatologia Brasileira**, Brasília, DF, v. 20, n. 2, p.184-189, mar. 1995.
- GODOY, C.V. **Influência de variáveis climáticas no desenvolvimento da ferrugem polysora na cultura do milho**. 2000. 79f. Dissertação (Doutorado em Fitopatologia) – Escola Superior de Agricultura “Luiz de Queiroz”. Piracicaba.
- GODOY, C.V. et al. Temporal progress of southern rust in maize under different environmental conditions. **Fitopatologia Brasileira**, Brasília, DF, v. 28, n.3, p.273-278, May/June 2003. <http://dx.doi.org/10.1590/S0100-41582003000300008>
- JULIATTI, F.C.; SANTOS, M.A. Métodos de avaliação de doenças de plantas. In: LUZ et al. **Revisão anual de patologia de plantas**. Passo Fundo, 1999. v. 7, p.407-454.
- McGEE, D.C. **Maize disease: a reference source for seed technologists**. Saint Paul: Ames: American Phytopathological Society, 1988. 150p. [http://dx.doi.org/10.1016/S0012-3692\(16\)30548-7](http://dx.doi.org/10.1016/S0012-3692(16)30548-7)
[http://dx.doi.org/10.1016/S0012-3692\(16\)33573-5](http://dx.doi.org/10.1016/S0012-3692(16)33573-5) [http://dx.doi.org/10.1016/S0012-3692\(16\)30546-3](http://dx.doi.org/10.1016/S0012-3692(16)30546-3)

MELCHING, J.S. Corn Rusts: types, races, and destructive potencial. In: ANNUAL CORN AND SORGHUM RESERCH CONFERENCE, 30., 1975. **Proceedings...** Washington: **American Seed Trade Association, 1975. p. 90-115.**

OLIVEIRA, E.B.M. **Seleção de genitores resistentes a *Puccinia polysora*, *Physopella zae* e *Phaeosphaeria maydis* em milho.** 2001. 79f. Dissertação (Mestrado em Genética e Melhoramento de Plantas) – Universidade Federal de Goiás, Goiânia.

PEREIRA, O. A. P. Doenças do milho (*Zea mays* L.). In: KIMATHI, H. et al. **Manual de fitopatologia.** 3. ed. São Paulo: Ceres, 1997. v. 2, p.539-555.

SCHIEBER, E. Distribución de *Puccinia polysora* y *Puccinia sorghi* en Africa y su patogenicidad sobre plasma de germen de maiz latinoamericano. **Boletín Fitossaniatario de La Fao**, St. Paul. v. 19, p.25-31, feb. 1971.

SHERFF, A.F.; MACNAB, A.A. **Vegetables diseases and their control:** corn. 2.ed. New York: John Wiley & Sons, 1986. p.232-34.

SHURTLEFF, M. C. **Compendium of corn diseases.** 2.ed. Saint Paul: American Phytopathological Press, 1992. 105p.

VIEIRA, R. A. et al. Genetic resistance of new popcorn hybrids to foliar diseases. **Crop Breeding and Applied Biotechnology**, Viçosa, v.9, n.,140146,2009.

VON PINHO R. G. et al. Reação de híbridos comerciais de milho às ferrugens polissora e tropical. **Pesquisa Agropecuária Brasileira Brasília**, v. 36, n. 3 p. 439-445, mar. 2001. <http://dx.doi.org/10.1590/S0100-204X2001000300007>

VON PINHO, R.G. et al. Comparação de métodos para quantificação da severidade das ferrugens polissora e tropical do milho. **Ciências Agrotécnicas**, Lavras, v.24, n.1, p.22-36, jan./mar., 2000.

WHITE, D. G. **Compendium of corn diseases.** 3. ed. St. Paul: American Phytopathological Society. 1999. 78p.