



SPECIFIC CHARACTER OF OBJECTIVE METHODS FOR DETERMINING WEIGHTS OF CRITERIA IN MCDM PROBLEMS: Entropy, CRITIC, SD

Irik Z. Mukhametzyanov^{1,2*}

¹ Ufa State Petroleum Technological University, Ufa, Russia

² Center for Strategic and Interdisciplinary Research, Ufa Federal Research Centre of
the Russian Academy of Sciences, Ufa, Russia

Received: 22 March 2021;

Accepted: 18 May 2021;

Available online: 11 June 2021.

Original scientific paper

Abstract. The comparative analysis of objective methods for determining the weights of criteria in the problems of multi-criteria decision-making is carried out. It is shown that the use of methods for determining the weights of criteria, based on formal processing of the decision matrix (Entropy, CRITIC, Standard deviation) for MCDM problems in some cases is not correct. It is demonstrated that the Entropy weighting method (EWM) is highly sensitive to evaluation of probabilities of states based on the decision matrix. For the Entropy method two modifications of estimation of probabilities of states are proposed that partially eliminate the contradictions of the basic EWM method. The first modification (EWM.df) is based on a statistical approach and it estimates the probabilities of states based on attribute distribution function. The second modification (EWM.dsp) estimates the probabilities of states based on the relative dispositions of attributes. Two options both have their supporting rationale. The analysis of integrated weighing methods is carried out and various options for aggregation of weights are given. An integrated EWM-Corr-method is proposed which allows to re-allocate the weights obtained by the Entropy method among correlated criteria.

Key words: *Multi-criteria decision making, weights estimation, Entropy weighting method, CRITIC method, Standard Deviation method, integrated weighting methods.*

* Corresponding author.

E-mail addresses: IZmukhametzyanov@gmail.com (I. Z. Mukhametzyanov).

1. Introduction

A significant number of multi-criteria decision-making methods (MCDM) constructs the performance indicator of alternatives, taking into account the weights of the criteria (Hwang & Yoon, 1981; Triantaphyllou, 2000; Tzeng & Huang, 2011; Pamucar & Savin, 2020). The weights of the criteria quantify their importance and can significantly affect the outcome of the decision-making process. Many authors note the high sensitivity of the solution to variations in the weights of the criteria (Barron & Schmidt, 1988; Mareschal, 1988; Wolters & Mareschal, 1995; Li et al., 2013; Karande et al. 2016; Mukhametzhanov & Pamučar, 2018; Zolfani et al., 2020). Given the importance of the weighing procedure, the balanced choice requires comprehensive analysis.

As you know, the weights of the components of a complex system can be obtained in different ways. One of the generally accepted classifications of methods for assessing criteria weights divides methods into three categories based on subjective, objective and integrated or combined approach to weighing (Goodwin & Wright, 1998, Bobko et al., 2007; Ginevicius & Podvezko, 2005; Jahan et al., 2012, Odu, 2019; Ali et al., 2021).

The determination of the subjective weight is based on the opinion of experts or expert groups representing the views of various stakeholders. These are such methods as the direct ranking method (DR) (Goodwin & Wright, 1998; Roberts & Goodwin, 2002; Von Winterfeldt & Edwards, 1986), the point allocation (PA) method (Doyle et al., 1997; Roberts & Goodwin, 2002), the ranking method (Ahn & Park, 2008; Barron, 1992, 1996; Roberts & Goodwin, 2002; Solymosi & Dombi, 1986; Milosevic et al., 2021), methods of programming (Pekelman & Sen, 1974; Shirland et al., 2003; Deng et al., 2004), Delphi method (Hwang & Yoon, 1981), pair-wise comparison (AHP) (Saaty, 1980; Takeda et al., 1987), step-wise weight assessment ratio analysis (SWARA) (Kersulienė et al., 2010), full consistency method (FUCOM) (Pamučar et al., 2018; Pamucar & Ecer, 2020), Level Based Weight Assessment (LBWA) (Žižović & Pamucar, 2019). One of the important problems of subjective methods is an assessment of the consistency of expert opinions. For example, the AHP-method defines a consistency index, which improves the reliability of the weight estimates. Other procedures for assessment of the consistency of expert judgments are based on statistical methods and correlation.

The category of objective assessment methods is based on the use of information about the criteria and their interactions contained in the decision-making matrix. These are such methods as entropy weighting method (EWM) (Lotfi & Fallahnejad, 2010; Wu et al., 2011; He et al., 2016), CRiteria Importance Through Inter-criteria Correlation (CRITIC) (Diakoulaki et al., 1995), standard deviation (SD) and their modifications (Jahan et al., 2012; Žižović et al., 2020; Alostia et al., 2021). For these methods, there is no answer to the question of how fully and objectively the limited sample of attributes of alternatives describes the value of the criteria. Obviously, the result is completely determined by the decision matrix. In particular, this article provides examples of how strongly the weights obtained for this category of methods for specially constructed decision-making matrices can differ. This paper analyzes the impact of normalization of the decision matrix and inversion of cost attributes on the weights estimation results. The hypersensitivity of the Entropy method to the estimation of state probabilities based on the decision matrix is

shown. For the Entropy method, a statistical approach to assessing the probability of states is proposed, which partially eliminates the contradictions of the formal method.

The essence of the integrated methods consists in combining the weighing results obtained by different methods with the subsequent use of the normalization of values by the Sum-method (Ma et al., 1999; Xu, 2004; Ustinovičius, 2001; Jahan et al., 2012; Vinogradova et al., 2018; Đalić et al., 2020; Singh et al., 2020). It is believed that integration can overcome the disadvantages of subjective and objective weighing methods. The integration of weights obtained by different methods takes different forms. In some cases, the combination procedure strengthens (weakens) the weights for “strong” (“weak”) criteria, when the criterion had a greater (lower) weight in both weighing methods, in other cases, the weight of the criteria is smoothed out when the criterion had a larger one in one of the methods and less weight in a different weighing method.

The paper presents various options for aggregating weights. Numerical examples show the importance of preliminary analysis of the results when choosing the final version of the aggregation procedure, taking into account the peculiarities of the decision-making problem. An integrated EWM-Corr method is proposed, which allows redistributing the weights obtained in the Entropy method between correlated criteria.

The idea that the assessment of weights is purely technical in nature and therefore objective is erroneous. It is rather difficult to make them objective, since there can be different problems with different data structures, varying degrees of uncertainty, etc. The existence of many methods and their modifications leading to different results indicates that the estimation of weights requires serious analysis. Since the design and determination of weights can always be interpreted in terms of value judgments, the procedure should also include the subjective opinions of individual experts. The differences in assessments of the surrounding world, as noted by the famous subjective idealist George Berkeley, are due to the fact that people's perceptions are far from identical. Their correct agreement limits the decision obtained by majority rule.

The purpose of this work is to show the characteristic features, shortcomings and possible contradictions of some methods for evaluating the weights of criteria for complex and critical analysis when solving problems of making multi-criteria decisions.

2. Preliminaries. Objective weighting methods: Entropy, CRITIC, SD

The category of objective assessment methods is based on the use of information about the criteria and their interaction contained in the decision-making matrix. The standard deviation or an “entropy” measure of importance have been proposed for quantifying contrast intensity and thus deriving objective weights of criteria (Zeleny, 1982). For objective weight estimation methods, it is believed that the greater the scatter of the values of the attribute of alternatives (the greater the difference between the values of the elements in the column, or the greater the variance, or the less entropy), the greater valuable information the criterion (indicator) contains and the higher the criterion weight. The criterion in which all alternatives have the same performance does not offer any additional information. The general structure of

Specific character of objective methods for determining weights of criteria...

objective methods for assessing weights is shown in the scheme in Figure 1. Data transformation (or normalization) is intended to co-ordinate the measurement scales of individual attributes. In order to maintain the proportions between natural and normalized values, linear transformations are used.




Figure 1. General structure of objective methods for evaluating the weight of criteria.

Entropy — EWM-method, multiplication of standard deviation and correlation — CRITIC-method and standard deviation — SD-method are used as key indicators. The weights of the criteria are determined as the intensities of the key indicators using the Sum-normalization.

Below are step-by-step weight estimation algorithms using the three most commonly used methods for solving MCDM problems — Entropy, CRITIC and SD.

Nomenclature:

- A_i alternatives (objects) ($i=1, \dots, m$)
- C_j^+, C_j^- criteria or objects properties ($j=1, \dots, n$), (+)benefit, (-) cost
- a_{ij} elements of decision matrix (DM)
- r_{ij} normalized elements of decision matrix
- \bar{r}_j average value of j -th criterion
- a_j^{\max} maximum element in criteria j
- a_j^{\min} minimum element in criteria j
- w_j weight or importance of criteria ($j=1, \dots, n$)

2.1. Entropy weighting method (EWM)

The values of the decision matrix are transformed into the segment [0; 1] using Max-Min normalization (1) with simultaneous inversion (2) of cost criteria values:

$$r_{ij} = \frac{a_{ij} - a_j^{\min}}{a_j^{\max} - a_j^{\min}}, r_{ij} \in [0; 1] \quad (1)$$

$$r_{ij} = \frac{a_j^{\max} - a_{ij}}{a_j^{\max} - a_j^{\min}}, r_{ij} \in [0; 1] \quad (2)$$

The intensity (p_{ij}) of the j -th attribute of the i -th alternative is calculated for each criterion (Sum-method):

$$p_{ij} = \frac{r_{ij}}{\sum_{i=1}^m r_{ij}}, \forall i = 1, \dots, m, j = 1, \dots, n; \sum_{i=1}^m p_{ij} = 1 \tag{3}$$

To calculate the entropy (e_j) and the key indicator (q_j) of each criterion:

$$e_j = -\frac{1}{\ln m} \cdot \sum_{i=1}^m p_{ij} \cdot \ln p_{ij}, j=1, \dots, n; \text{ (if } p_{ij}=0 \Rightarrow p_{ij} \cdot \ln p_{ij}=0) \tag{4}$$

$$q_j = 1 - e_j, j = 1, \dots, n \tag{5}$$

To calculate the weight of each criterion:

$$w_j = q_j / \sum_{k=1}^n q_k, j = 1, \dots, n \tag{6}$$

The entropy of the attributes of alternatives for each criterion is a measure of the significance of this criterion. It is believed that the lower the entropy of the criterion, the more valuable information the criterion contains.

2.2. CRiteria Importance Through Inter-criteria Correlation (CRITIC)

The values of the decision matrix are transformed based on the concept of the ideal point. To determine “best” ($B=b_j$) and “worst” ($T=t_j$) solution ($[1 \times n]$ -vector) for all attributes and determine relative deviation matrix $V[m \times n]$:

$$r_{ij} = \frac{(a_{ij} - b_j)}{(b_j - t_j)} \tag{7}$$

To determine standard deviation (s) ($[1 \times n]$ -vector) for colls of V :

$$s_j = \text{std}(V) = \sqrt{\frac{1}{m-1} \cdot \sum_{i=1}^m (r_{ij} - \bar{r}_j)^2} \tag{8}$$

To determine the linear correlation matrix (c_{jk}) ($[n \times n]$ -matrix) for colls of V is the (correlation coefficient between the vectors r_j and r_k):

$$c_{jk} = \text{corr}(V) = \frac{\sum_{i=1}^m (r_{ij} - \bar{r}_j)(r_{ik} - \bar{r}_k)}{\sqrt{\sum_{i=1}^m (r_{ij} - \bar{r}_j)^2 \sum_{i=1}^m (r_{ik} - \bar{r}_k)^2}}, j, k = 1, \dots, n \tag{9}$$

To calculate the key indicator and weight of criteria by the formula (6):

$$q_j = s_j \cdot \sum_{k=1}^n (1 - |c_{jk}|), j = 1, \dots, n \tag{10}$$

In the CRITIC method, the standard deviation s_j is a measure of the significance of this criterion. Allowance for the relationship between the criteria is determined through the correlation matrix, which allows you to distribute the weight between the correlated criteria through the coefficients of reduction $(1-c)$. The amount

Specific character of objective methods for determining weights of criteria...

shown in expression (10) is a measure of the conflict created by j -th criterion in relation to the rest of the criteria. Finally, the amount of information contained in the j -th criterion is determined using multiplicative aggregation of measures by the formula (10).

The Spearman rank correlation coefficient could be used instead of c_{jk} , in order to provide a more general measure of the relationship between the rank orders of the elements included in the vectors r_j and r_k .

2.3. Standard deviation (SD)

The values of the decision matrix are transformed into the segment $[0; 1]$ using Max-Min normalization with simultaneous inversion of cost criteria values by formulas (1) and (2).

To calculate the key indicator and weight of criteria by expression (6):

$$q_j = s_j = \text{std}(r_{ij}) = \sqrt{\frac{1}{m-1} \cdot \sum_{i=1}^m (r_{ij} - \bar{r}_j)^2} \quad (11)$$

The standard deviation of r_j is a measure of the value of that criterion to the decision making process.

3. Features of methods for evaluating the weights of criteria in MCDM problems

3.1. Estimating the probabilities of the object states in the Entropy method

One of the main notes for the Entropy method is that the p_{ij} in the entropy method determines the probabilities of m possible and independent states (alternatives) of an object. It is not clear how the probabilities of states and intensities, obtained by the Sum-method according to expression (3), are related.

If we combine expressions (1)–(3) for profit criteria (similarly for cost criteria), we get:

$$p_{ij} = \frac{a_{ij} - a_j^{\min}}{\sum_i (a_{ij} - a_j^{\min})} \quad (12)$$

The p_{ij} values are the relative distances of the j -th attribute of alternatives to the alternative with the smallest value.

Obviously, a discrete random variable is determined by a pair (x_i, p_i) and the values of p_i are needed to calculate the entropy. However, expression (3) determines p_i with respect to x_i . In Sections 4.2 and 4.3, to determine the probabilities of possible states of alternatives, it is proposed to use the distribution functions of the attributes of alternatives built on the values of the decision matrix and to use relative dispositions of attributes instead of p_{ij} .

3.2. Difference in weights of criteria when changing many alternatives

Consider a decision-making problem in which some or all of the set of alternatives are different, and the attributes are assessed within the same criteria. Taking the principle of uniformity and continuity of changes in the properties of an object, such groups of alternatives exist and have attributes from a certain interval of the area of change. Table 1 presents decision matrices for two groups of alternatives to one problem in the context of the same criteria.

Let's form a mix-matrix of solutions *DM-3* obtained by replacing the first alternative *A*₁ of the matrix *DM-1* with the first alternative *B*₁ from *DM-2*, and form more one decision matrix *DM-12* of dimension [16x5], obtained by the concatenation of the matrices *DM-1* and *DM-2*. Both sets of alternatives *A*_{*i*} and *B*_{*i*} are related to the context of the problem: alternatives are objects of the same nature,

Table 1. Decision matrices *DM-1* and *DM-2* for two groups of alternatives *A*_{*i*} and *B*_{*i*} in the context of the same criteria.

		<i>DM-1</i> , criteria: benefit(+)/cost(-)					<i>DM-2</i> , criteria: benefit(+)/cost(-)					
		<i>C</i> ₁₊	<i>C</i> ₂₋	<i>C</i> ₃₊	<i>C</i> ₄₊	<i>C</i> ₅₋	<i>C</i> ₁₊	<i>C</i> ₂₋	<i>C</i> ₃₊	<i>C</i> ₄₊	<i>C</i> ₅₋	
Alternatives	<i>A</i> ₁	71	4500	150	1056	478	<i>B</i> ₁	83	5322	170	1682	500
	<i>A</i> ₂	85	5800	145	2680	564	<i>B</i> ₂	84	6021	155	2140	513
	<i>A</i> ₃	76	5600	135	1230	620	<i>B</i> ₃	76	4219	155	1613	454
	<i>A</i> ₄	74	4200	160	1480	448	<i>B</i> ₄	73	6154	157	2047	582
	<i>A</i> ₅	82	6200	183	1350	615	<i>B</i> ₅	78	5453	173	2136	598
	<i>A</i> ₆	81	6000	178	2065	580	<i>B</i> ₆	82	6030	174	1238	587
	<i>A</i> ₇	80	5900	160	1650	610	<i>B</i> ₇	80	4344	172	1365	507
	<i>A</i> ₈	85	6500	140	1650	667	<i>B</i> ₈	77	6114	158	1585	592
std		5.1	813	17.3	518	74.6		3.8	788	8.7	347	54.5

having similar properties within the same criteria and having approximately the same range of variation. It should be expected that the criterion weights calculated by the same method for different decision matrices will be approximately the same. However, the weights of the criteria (and the ranks of alternatives) differ significantly (Table 2) not only for different methods, but within the same method for a different set of alternatives.

To characterize the deviation of the weights obtained in various examples and methods, the relative error is used (the calculation is made for each *j*-th components) $\delta_{s,k} = |w^{(s)} - w^{(k)}| / w^{(s)} \cdot 100, \%$.

Table 2. Weights of the criteria for the decision matrix *DM-1* and *DM-2* of Table 1, the mix-matrix *DM-3*, and the concatenation matrix *DM-12*.

Entropy	<i>w</i> ₁	<i>w</i> ₂	<i>w</i> ₃	<i>w</i> ₄	<i>w</i> ₅	Rank (SAW)
<i>DM-1</i>	0.146	0.228	0.211	0.223	0.191	A2>A4>A6>...
<i>DM-2</i>	0.107	0.287	0.240	0.134	0.232	B3>B7>B1>...
<i>DM-3</i>	0.144	0.216	0.200	0.249	0.191	A2>A4>B1>...
<i>DM-12</i>	0.159	0.280	0.165	0.220	0.176	B3>A4>A2>...
mean	0.139	0.253	0.204	0.207	0.198	
std	0.022	0.036	0.031	0.050	0.024	
CRITIC						
<i>DM-1</i>	0.144	0.147	0.322	0.214	0.173	A2>A6>A4>...
<i>DM-2</i>	0.194	0.199	0.229	0.203	0.176	B3>B5>B1>...
<i>DM-3</i>	0.179	0.147	0.292	0.219	0.164	A2>A6>A1>...
<i>DM-12</i>	0.187	0.184	0.254	0.193	0.182	A2>B3>B1>...
mean	0.176	0.169	0.274	0.207	0.174	

Specific character of objective methods for determining weights of criteria...

std	0.022	0.026	0.041	0.012	0.007	
SD						
DM-1	0.210	0.203	0.207	0.183	0.196	A2>A4>A6>...
DM-2	0.175	0.206	0.231	0.195	0.192	B3>B1>B5>...
DM-3	0.216	0.181	0.220	0.189	0.193	A2>B1>A4>...
DM-12	0.208	0.225	0.192	0.176	0.199	A2>B3>A4>...
mean	0.202	0.204	0.213	0.186	0.195	
std	0.018	0.018	0.017	0.008	0.003	
	relative error of means δ_{s-k} , %					
Entropy-CRITIC	19.1	40.9	15.1	31.4	36.2	
CRITIC-SD	79.7	36.4	23.1	31.0	54.3	
Entropy-SD	45.3	19.4	4.4	10.1	1.5	

Table 2. (Continued)

DM-1	w_1	w_2	w_3	w_4	w_5	Rank (SAW)
Entropy	0.146	0.228	0.211	0.223	0.191	A2>A4>A6>...
CRITIC	0.144	0.147	0.322	0.214	0.173	A2>A6>A4>...
SD	0.210	0.203	0.207	0.183	0.196	A2>A4>A6>...
DM-2						
Entropy	0.107	0.287	0.240	0.134	0.232	B3>B7>B1>...
CRITIC	0.194	0.199	0.229	0.203	0.176	B3>B5>B1>...
SD	0.175	0.206	0.231	0.195	0.192	B3>B1>B5>...
DM3						
Entropy	0.144	0.216	0.200	0.249	0.191	A2>A4>B1>...
CRITIC	0.179	0.147	0.292	0.219	0.164	A2>A6>A1>...
SD	0.216	0.181	0.220	0.189	0.193	A2>B1>A4>...
DM-12						
Entropy	0.159	0.280	0.165	0.220	0.176	B3>A4>A2>...
CRITIC	0.187	0.184	0.254	0.193	0.182	A2>B3>B1>...
SD	0.208	0.225	0.192	0.176	0.199	A2>B3>A4>...

In this particular example, the largest variation in weights in the Entropy method is 185%, in the CRITIC method — 52% and SD — 24%. Such a wide spread leads to a change in the ranking of alternatives (Simple Additive Weighting method — SAW). On the one hand, all three matrices represent assessments of alternatives within the same criteria. However, the relative difference in the estimates of the weights for the given example for some criteria is up to 50% and more. Even when replacing one alternative, the weights of the criteria change significantly.

3.3. Decision matrix transformations and influence on key parameters of CRITIC and SD methods

Since key indicators (q) should not depend on scales of measurements of the attributes, the attributes should be normalized or reduced to dimensionless values. Considering that the number of methods of normalization and inversion totals about 20 (Jahan et al., 2015), we should expect the ambiguity of the results. 6 basic linear methods of normalization, presented in Table 3, have a meaningful interpretation. The IZ-method (Mukhametzyanov, 2018, 2019, 2021) is a generalization of the Max-Min method and will be used below in the Entropy-method.

Table 3. Basic linear methods for the normalization of the decision matrix.

non displacement: $r_{ij} = a_{ij} / k_j$			with displacement: $r_{ij} = (a_{ij} - a_j^*) / k_j$		
Max ^{*)}	Sum	Vec	Max-Min	dSum	IZ
$k_j = a_j^{\max}$; $k_j = \sum_{i=1}^m a_{ij} $; $k_j = \sqrt{\sum_{i=1}^m a_{ij}^2}$			$k_j = a_j^{\max} - a_j^{\min}$; $a_j^* = a_j^{\min}$.	$k_j = \sum_{i=1}^m (a_j^{\max} - a_{ij})$; $a_j^* = a_j^{\max} - k_j$.	$k_j = \frac{Z - I}{a_j^{\max} - a_j^{\min}}$,

^{*)} The short name of the normalization methods is determined by the semantic value of the compression ratio k . The method abbreviation is also used as the name of a function that converts values in accordance with the normalization method. For example, $r_{ij} = \text{Max}(a_{ij}) = a_{ij} / a_j^{\max}$. dSum-method is a combination of Max-Min and Sum methods (see the formula (12)) and 1-r inversion. In the IZ-method, I and Z define a segment of normalized values that is fixed for all criteria ($0 \leq I < Z \leq 1$).

The use of linear normalization methods is due to the fact that they retain the relative dispositions between the natural and normalized values of the alternatives:

$$\text{if } r_{ij} = \frac{a_{ij} - a_j^*}{k_j} \Rightarrow \frac{r_{pj} - r_{qj}}{\text{rng}(r_j)} = \frac{a_{pj} - a_{qj}}{\text{rng}(a_j)}, \forall p, q = \overline{1, m}, \tag{13}$$

where $\text{rng}(r_j) = r_j^{\max} - r_j^{\min}$ is the range of values for each attribute.

Preserving the “dispositions” of alternatives after normalization means that the normalized values preserve information about the system. For linear normalization without displacement, one degree of freedom is lost, and for normalization with displacement, two degrees of freedom are lost. Any nonlinear normalization leads to information distortion.

When using a general linear transformation of natural values $r_{ij} = (a_{ij} - a_j^*) / k_j$, the standard deviation is scaled:

$$s_j(r_{ij}) = \frac{s_j(a_{ij})}{k_j}, \tag{14}$$

and the matrix of pair-wise correlations does not change (invariant):

$$\text{corr}(r_{ij}) = \text{corr}(a_{ij}). \tag{15}$$

This means that the result of CRITIC and SD depends on the choice of the decision matrix normalization method, since the proportions between k_j in different normalization methods for different criteria will be different, and the intensity of the q_j key indicators will be different.

Figure 2 shows the domains of the normalized values of the attributes of the alternatives and shows the intervals $(m - s_j; m + s_j)$ of deviation from the mean m . For normalization methods, except for Max-Min and IZ, domains for different criteria have a different range of values $\text{rng}(r_j)$. The standard deviation s correlates with the range of domain sizes.

Specific character of objective methods for determining weights of criteria...




Figure 2. Domains of normalized values of attributes of alternatives and intervals ($m-s$; $m+s$) for various transformations of the decision matrix $DM-1$.

Of the six normalization options in Table 3, the Max-Min and IZ method is preferable, since in this case the range of values of all attributes is the same and the standard deviation for the normalized values $s_j(a_{ij})/k_j$ of various attributes does not depend on the range and has a natural interpretation as the degree of dispersion of values. In all other cases of normalization, the key indicators $q_j = s_j$ for various criteria are not consistent with each other, and the result depends on the transformation method of criteria and measurement scales. The numerical results of the example under consideration are shown in Table 4.

The results demonstrate a significant difference in criterion weights for different methods, which indicates the importance of choosing the correct normalization method. For the Max-Min and IZ normalization methods, the standard deviation is proportional to the factor $(Z-I)$. Therefore, the criteria weights will be equal (highlighted in the table).

Table 4. The values of the standard deviation and weights of the criteria of the decision matrix $DM-1$ for various transformation methods.

Norm. method	Standard deviation					Weights of criteria				
	C_{1+}	C_{2-}	C_{3+}	C_{4+}	C_{5-}	w_1	w_2	w_3	w_4	w_5
Max	0.060	0.125	0.095	0.193	0.112	0.103	0.214	0.162	0.331	0.191
Sum	0.008	0.018	0.014	0.039	0.016	0.084	0.190	0.144	0.411	0.170
Vec	0.023	0.051	0.039	0.107	0.046	0.086	0.192	0.147	0.403	0.172
Max-Min	0.366	0.354	0.361	0.319	0.341	0.210	0.203	0.207	0.183	0.196
dSum	0.111	0.111	0.081	0.063	0.099	0.239	0.239	0.175	0.135	0.213
IZ(0.2; 0.7)	0.183	0.177	0.180	0.160	0.170	0.210	0.203	0.207	0.183	0.196

For the CRITIC method, the general transformation (7), in contrast to the Max-Min normalization by the formula (1) is not correct, because the “best” (B) and “worst” (T) solution may differ from the maximum and/or minimum value. For example, if “best” and “worst” are absolute (from the entire set of possible alternatives, and not from a private sample in the investigated decision-making problem), then the normalized values are within the interval (0; 1), and for each criterion the domain

range will be different. The normalized values will be offset relative to the segment [0; 1] with all the consequences of changing the standard deviation of the attributes of individual criteria and changing their weight. Therefore, the correct application of the CRITIC method is possible only with the Max-Min transformation.

To invert the attributes of cost criteria, it is recommended the ReS-algorithm in two forms (Mukhametzhanov, 2020):

1) for natural values of attributes

$$a_{ij^*} = -a_{ij^*} + a_{j^*}^{\max} + a_{j^*}^{\min}, \forall j^* \in C_j^- \tag{16}$$

2) for the normalized values of attributes

$$\begin{aligned} 1) & r_{ij} = Norm(a_{ij}), \forall j = 1, \dots, n \\ 2) & \tilde{r}_{ij^*} = -r_{ij^*} + r_{j^*}^{\max} + r_{j^*}^{\min}, \forall j^* \in C_j^- \end{aligned} \tag{17}$$

where $Norm()$ is one of the linear normalization methods applied to both benefit and cost attributes; the j^* index meets the cost criteria. The abbreviation of the normalization method (Table 3) or inversion is also used as the name of the function that transforms values in accordance with the selected method: $r_{ij} = Vec(r_{ij}), v_{ij} = ReS(r_{ij})$. ReS-algorithm is universal for any linear and non-linear transformations and ReS-inversion algorithm preserves dispositions of natural and normalized attribute values. For the Max-Min normalization method, the inversion by the expression (2) and the inversion using the ReS-algorithm have the same results.

According to the standard deviation expression, inverting cost attributes does not affect the outcome.

However, in the matrix of pair-wise correlations, the sign of some of the coefficients changes according to the rule:

$$corr(ReS(r_{ij})) = corr(r_{ij}) \otimes \overline{g}^T \cdot \overline{g} \tag{18}$$

where \otimes is the element-wise product of two matrices of dimension $[n \times n]$;

$\overline{g} = (\text{sgn } C_1, \text{sgn } C_2, \dots, \text{sgn } C_n)$ $[1 \times n]$ -vector

$$\text{sgn } C_j = \begin{cases} +1, & \text{if } C_j - \text{benefit criteria} \\ -1, & \text{if } C_j - \text{cost criteria} \end{cases}$$

For example, for the test matrix $DM-1$ used in this paper:

$$\overline{g}^T \cdot \overline{g} = \begin{pmatrix} 1 \\ -1 \\ 1 \\ 1 \\ -1 \end{pmatrix} \cdot (1 \ -1 \ 1 \ 1 \ -1) = \begin{pmatrix} 1 & -1 & 1 & 1 & -1 \\ -1 & 1 & -1 & -1 & 1 \\ 1 & -1 & 1 & 1 & -1 \\ 1 & -1 & 1 & 1 & -1 \\ -1 & 1 & -1 & -1 & 1 \end{pmatrix}$$

Negative correlation should not increase the weight of the key factor in expression (10). Therefore, in expression (10), the correlation must be absolute and this is the correct expression for CRITIC.

Specific character of objective methods for determining weights of criteria...

3.4. How much do the weights of the Entropy method vary depending on the distribution in the data?

To answer this question, let us turn to the graphical illustration (Fig. 3), which shows the function $f(x) = -x \cdot \ln x$; the values p_{ij} of attributes of each alternative after transformation Sum; the values of the components of the entropy of the j -th criterion $e_{ij} = -p_{ij} \cdot \ln p_{ij}$ of each alternative; and domains of values of e_{ij} for each criterion, allowing to observe the nature of the distribution. The values $-p_{ij} \cdot \ln p_{ij}$ for all j are actually on the line of the function $-x \cdot \ln x$ and, for clarity the images, are separated by a parallel translation along the OY axis.

The entropy function $f(x)$ has a maximum at the point $1/e \approx 0.368$. For m states, the probabilities p_i have, on average, less value than $1/e$, and are concentrated in the vicinity of the point $1/m$. Accordingly, the entropy component $-p_i \cdot \ln p_i$ is concentrated in the vicinity of the point $1/m \cdot \ln(1/m)$.

The values of the weight of the criterion and the entropy of the criterion are opposite. In accordance to the expression (4), the maximum value of the criterion entropy is equal to 1, and the criterion weight is equal to 0. This state is achieved when all probabilities $p_{ij} = 1/m$ for a fixed j .




Figure 3. Entropy and entropy components $(-x \cdot \ln x)$ for the DM-1 matrix.

This is possible provided that all alternatives have the same value for the j -th attribute $p_{ij} = Const$. The greater the difference between the values of the j -th attribute for various alternatives, the lower the value of the criterion entropy and, in accordance to the expression (5), the greater the value of the criterion weight. This is due to the specifics of data normalization by the Sum-method.

Therefore, the criterion j , the attributes of the alternatives of which are concentrated closer to the smallest value a_j^{\min} , will have more weight. This is believed to indicate that the criterion contains valuable information. A very controversial assumption. The weak point of the entropy method for assessing weight is the high sensitivity (hypersensitivity) of weight to the values of the entropy of various criteria. This hypersensitivity is due to the exponential behavior of the logarithm in the vicinity of 0. Indeed, the rate of change of entropy along the i -th component is determined by expression:

$$\frac{\partial e_j}{\partial p_i} = -\frac{1}{\ln m} \cdot (\ln p_{ij} + 1) \tag{19}$$

therefore, small changes in the values of the j -th attribute of the i -th alternative can change the weights. For example, (see Table 2 and Fig. 3) the entropies for criteria 2 and 3 differ by 1.5%, while the weights differ by 8%. A slight error in the estimates of the attribute of alternatives will lead to a significant change in the estimates of the weights.

Another weak point of the entropy method is the sensitivity to the distribution of data in the domain of p_{ij} values. The concentration of the normalized values of one of the attributes in the vicinity of 0 sharply decreases the entropy and leads to an increase in the weight of this criterion. Figure 4 illustrates this feature. When replacing only one element a_{12} in the decision matrix $DM-1$ from 85 to 72, entropy of the first criterion decreased from 0.8897 to 0.8065, and the weight of the first criterion increased from 0.146 to 0.231 (by 158%). The ranking of alternatives has also changed.




Figure 4. Weights of criteria and entropy with decreasing variance of the 1st attribute in the decision matrix $DM-1$.

Note that when the data changed, the standard deviation for the first criterion increased from 5.120 to 5.125 (natural). The SD-method shows weight changes of only 0.1% — from 0.2102 to 0.2104. This example demonstrates the high sensitivity (hypersensitivity) of EWM-method to the distribution of data in the domain.

Specific character of objective methods for determining weights of criteria...

The apparent efficiency of the entropy method can most likely be attributed to a situation in which the attributes of the alternatives have approximately the same distribution and approximately the same variance. In such a situation, the weights of the criteria differ less significantly.

3.5. Do you need to transform (normalize) the natural values of the attributes before applying the Entropy method?

Some studies when estimating weights do not use preliminary data transformation (by formulas (1) and (2)), believing that it is obvious that the transformation Sum (formula (3)) is sufficient, or they use normalization methods other than Max-Min. For example, in (Wu et al., 2011; Li et al., 2011), values are normalized and inverted for cost criteria using the Max and Vec methods.

The peculiarity of the transformation according to the expression expression (3) (Sum-method) is that the normalized p_{ij} values are converted into the interval (0; 1) with the preservation of the dispositions of natural values, but the domains for different criteria have different sizes and are shifted relative to each other. This is clearly seen in the diagrams for p_{ij} in Figure 5. For more details on the displacement of domains during normalization, see the study by Mukhametzyanov (2020, 2021). The displacement of the p_{ij} values by the j -th criterion, according to the analysis carried out above in Section 3.4, strongly affects the entropy of the e_j criteria, which is reflected in the weights.

Normalization based on linear methods *without displacement* (Max, Sum, Vec) fully preserves the proportions of natural and normalized values of attributes. However, after the re-normalization, the equalities are valid (Mukhametzyanov, 2020):

$$p_{ij} = \text{Sum}(a_{ij}/k_j) = \text{Sum}(\text{Max}(a_{ij})) = \text{Sum}(\text{Sum}(a_{ij})) = \text{Sum}(\text{Vec}(a_{ij})) = \text{Sum}(a_{ij}). \quad (20)$$




Figure 5. Probabilities (p) of possible states of alternatives and entropy components ($-p \cdot \ln p$) of the Entropy method for the $DM-1$ matrix and the normalized decision matrix $DM-1$ by the Max-Min and dSum methods. Two-axis technique.

Therefore, the result of calculating the weights of the criteria will not change. In Figure 4, in particular, p_{ij} coincides for the case when the preliminary data transformation is not performed and for the cases of linear transformation without displacement (1 fragment).

Normalization based on linear methods with *displacement* (Max-Min, dSum) retains only the dispositions of the values. However the probabilities change:

$$p_{ij} = \text{Sum}((a_{ij} - a_j^*) / k_j) \# \text{Sum}(a_{ij}), \tag{21}$$

and therefore the entropies of the criteria will change.

In a significant part of the studies (Liang et al., 2006; Dong et al., 2016; Zhao et al., 2017; Chen et al., 2019; Wu et al., 2018), the entropy method uses preliminary transformation of the initial data using normalization of Max-Min with simultaneous inversion of the values of the cost criteria.

The Max-Min transformation maps attribute values of all criteria to the interval [0; 1], which aligns the domains for all criteria and eliminates their relative displacement. The Max-Min and inverse transformation for Max-Min also preserve the relative proportions of the values.

The subsequent transformation by the Sum-method leads to different values of the intensities (probabilities) p_{ij} than for the natural data of the decision matrix. At least one of the values after transformation by the Sum method is equal to 0. The transformed system is not identical to the original one. Despite the fact that agreement of domains for different criteria has been achieved, within the domain there is a shift in data towards a decrease in the contribution of individual attributes to the entropy of the criterion and a decrease in the entropy of the criterion. At least one of the values after the stepwise transformation by the Max-Min and Sum method is equal to 0. The transformations are uneven for various criteria (except for the distribution in the data that is symmetric with respect to transformation). Therefore, the previously transformed system is a different system, not identical to the original one, and the results of the weight estimation differ significantly.

Table 5 and Figure 4 shows an example of such calculations for the first criterion of the *DM-1* matrix.

Table 5. Probabilities of states of alternatives (p), entropy (e) and weights of criterion (w) for matrix *DM-1* and normalized matrix of solutions *DM-1* by Max-Min and dSum methods. EWM-method.

	p_{i1}			transform	e_1	e_2	e_3	e_4	e_5
	<i>DM-1</i>	Max-Min	dSum						
A_1	0.112	0	0.099	<i>DM-1</i>	0.9991	0.9949	0.9974	0.9804	0.9961
A_2	0.134	0.212	0.143	Max-Min	0.8897	0.8280	0.8410	0.8323	0.8558
A_3	0.120	0.076	0.115	dSum	0.9965	0.9961	0.9982	0.9989	0.9971
A_4	0.117	0.046	0.109		w_1	w_2	w_3	w_4	w_5
A_5	0.129	0.167	0.134	(1) <i>DM-1</i>	0.028	0.160	0.079	0.612	0.121
A_6	0.129	0.152	0.130	(2) Max-Min	0.146	0.228	0.211	0.223	0.191
A_7	0.126	0.136	0.127	(3) dSum	0.265	0.295	0.137	0.081	0.222
A_8	0.134	0.212	0.143	δ_{1-2}	421.4	42.5	167.1	63.6	57.9
				δ_{2-3}	81.5	29.4	35.1	63.7	16.2

The values of the entropy and weights of the criteria calculated after performing the preliminary data transformation can significantly differ from the analogous

Specific character of objective methods for determining weights of criteria...

values without transformation. The largest variation in weights in the Entropy method is over 400%, with no strong case for data transformation.

3.6. Do you need to apply value inversion for cost criteria before applying Entropy?

The answer to this question is not obvious. On the one hand, the criteria are independent and the entropy (e_j) of the criterion is not related to the direction of improving the values of alternatives. On the other hand, attribute values are ordered from lowest to highest value. When the direction of the coordinate axis and the origin is changed, the state probability distribution function is reflected. If the distribution does not have symmetry with respect to such a transformation, then formally the entropy of the system will change due to the relative displacement of values within the domain (even in the case of using the best of the inversion algorithms — ReS (Mukhametzyanov, 2020), which preserves the positions of the boundaries of the domain of inverted values). For example, for the example considered in Table 2, the weights calculated without preliminary inversion of the values of the cost criteria (Table 6) will differ from the weights calculated with preliminary inversion.

Table 6. Weights and ranks of decision matrices, calculated without preliminary inversion of the values of the cost criteria. EWM-method.

	w_1	w_2	w_3	w_4	w_5		w_1	w_2	w_3	w_4	w_5
<i>DM-1</i>	0.028	0.148	0.081	0.625	0.117	<i>DM-2</i>	0.030	0.278	0.036	0.525	0.132
rank	$A_2 > A_6 > A_4 > \dots$ (SAW)					rank	$B_2 > B_5 > B_4 > \dots$ (SAW)				

Thus, the second argument is strongly pointing in favor of pre-inverting the cost attribute values. For the inverse of values of cost criteria, the ReS-algorithm presented in 3.3 above is recommended.

It is not desirable to use non-linear inversion. For example, in the works of Li, et al., (2011), Wu, et al., (2011), a nonlinear transformation $iMax = a_{ij}^{*min} / a_{ij}^*$, is used, which transforms the original system to another system that is not equivalent to the original one, the weights of which are different.

3.7. IZ-transformation of natural values of attributes for the Entropy method

Applying linear operations of shift to a fixed point, tension-compression and subsequent displacement, it is possible to transform the natural values of all criteria into a fixed (or predetermined) interval of values $[a; b]$ (Mukhametzyanov, 2018, 2019, 2021) while maintaining the relative dispositions of natural values for each attribute. A special case of such a transformation is the Max-Min normalization method, which converts the attribute values for all criteria into the segment $[0; 1]$.

An example of IZ-transformation of the matrix of solutions *DM-1* into the segment $[0.2; 0.7]$ is shown above in Figure 2. IZ-transformation is in no way inferior, and even more correct than transformation in $[0; 1]$, which has problems with null values. For example, there is no contribution of min-value of attributes to entropy. As a result of the IZ-transformation, in comparison with the Max-Min transformation, the standard deviation and the matrix of pairwise correlations will not change. This means that such data transformation does not affect the result of the CRITIC and SD methods. However, p_{ij} will change, and due to the nonlinearity of

the entropy components $(-p_{ij} \cdot \ln p_{ij})$, both the entropy and the weights of the criteria will change.

It is natural to ask how the choice of the IZ-transformation interval $[a; b]$ affects the entropy and weights of the criteria?

Due to the strict monotonicity of the entropy components on the interval $[0, 1/e]$ (see Fig. 3), all changes will also be strictly monotonic, except for the cases when any of the values of p_{ij} are greater than $1/e$. Figure 6 shows a color map of entropy values




Figure 6. Color map of entropy values for the 5th criterion of $DM-1$ depending on the size and position of the transformation interval $[a; b]$ with respect to $[0; 1]$.

for the 5th criterion (the results are similar for other criteria) depending on the size and position of the transformation interval $[a; b]$ with respect to $[0; 1]$. According to such a map, entropy changes nonlinearly and strictly monotonically depending on changes in the size and position of the domain of normalized values.

Non-linear changes in entropy leads to the fact that the dynamics of the weight coefficients, firstly, is also non-linear (Figure 7). Secondly, there are points of change in the priority of criteria (intersection points), for which the priority of the criterion (its weight) changes. This indicates the dependence of the weighing result on the choice of the size and position of the domain.

Specific character of objective methods for determining weights of criteria...




Figure 7. Dynamics of entropy and weight coefficients depending on the size and position of the domain $[a; b]$ at IZ-transformation (correspondence according to the Figure 6). *DM-1*.

The conclusion from the analysis is not encouraging. When manipulating data transformation in the Entropy-method, you can manipulate the criteria weights.

3.8. Integrated methods

The essence of integrated methods (or aggregation methods) is to combine weighing results obtained by different methods. For example, weights obtained using subjective and objective weighing methods such as AHP and Entropy are often corrected. It is assumed that the combined weighing method will reduce the potential bias of a single subjective or objective weight, or can make up for the deficiency of the subjective weight.

Let the weights $w_j^{(1)}$ of the j -th criterion be obtained by one of the available methods, and the weights $w_j^{(2)}$ be obtained by another method. A common aggregation procedure has the following multiplicative form:

$$q_j = w_j^{(1)} \cdot w_j^{(2)} \quad (22)$$

Aggregation (21) strengthens (weakens) the weights for “strong” (“weak”) criteria, when the criterion has a greater (lower) weight in both weighing methods. In other cases, the weight of the criteria is smoothed, when the criterion had greater weight

in one of the methods and less weight in the other weighing method. In some cases, such a procedure is dangerous from the point of view of amplifying the error.

The q_j value can be arranged in various forms, for example, as different variants of the average value in N weighing methods:

– harmonic mean (HM)

$$q_j = N / \sum_{k=1}^N \frac{1}{w_j^{(k)}} \quad (23)$$

– geometric mean (GM)

$$q_j = \left[\prod_{k=1}^N w_j^{(k)} \right]^{1/N} \quad (24)$$

– arithmetic mean (AM)

$$q_j = \frac{1}{N} \sum_{k=1}^N w_j^{(k)} \quad (25)$$

– root sum of squares (RSSq)

$$q_j = \left[\sum_{k=1}^N w_j^{(k)2} \right]^{-1/2} \quad (26)$$

The weights are calculated as the intensities of the integrated indicator q_j by formula (6):

$$w_j = q_j / \sum_{j=1}^n q_j \quad (27)$$

For different means, the following inequalities hold:

$$\min \leq \text{HM} \leq \text{GM} \leq \text{AM} \leq \max \leq \text{RSSq} \quad (28)$$

The choice of the method for the mean is not formalized and is determined by the context of the problem.

The weights aggregation method is compensating. It compensates for low criterion weights in some intermediate weighing methods with high values in others. If compensation is acceptable, then a higher weight value for the same criterion in a different weighing method will be required to obtain a higher weight value. With multiplicative aggregation, a higher value of the weight of one of the criteria can be obtained by reducing the weight of the other criterion.

Given the variability of aggregation and compensation, the integrated methods are not general. Rather, they give the researcher a formal opportunity to eliminate the discrepancy between the criteria weights in a particular problem.

Specific character of objective methods for determining weights of criteria...

4. EWM modification

4.1. Modification of the entropy method using correlation: EWM-Corr

As discussed in section 3.8 above, key indicators of different methods can be combined to assess criterion weights. The combination of entropy and correlation in the following multiplicative form is relevant:

$$q_j = (1 - e_j) \cdot \sum_{k=1}^n (1 - |c_{jk}|), \quad j = 1, \dots, n \quad (29)$$

where e_j is the entropy of the criterion determined by formulas (1)–(4), c_{jk} is the matrix of pairwise correlations determined by the formula (9).

The first factor determines the dispersion of the attribute values, and the second factor allows the weight to be distributed among the correlated criteria in terms of reduction factors $(1-c)$.

The weights of the criteria are determined by the intensity of the key indicator q_j using the Sum-normalization by the formula (6).

Table 7 shows the results of calculating the weights by the proposed EWM-Corr method.

Table 7. Weights of the criteria and rank for the decision matrix *DM-1*, *DM-2*, *DM-3*, *DM-12*, obtained using the EWM-Corr method.

Decision matrix	w_1	w_2	w_3	w_4	w_5	Rank (SAW)
DM-1	0.098	0.161	0.321	0.255	0.165	A2>A6>A4>...
DM-2	0.121	0.281	0.241	0.141	0.216	B3>B7>B1>...
DM-3	0.118	0.173	0.263	0.285	0.160	A2>A6>AB1>...
DM-12	0.145	0.230	0.219	0.244	0.163	A2>B3>B5>...
mean	0.139	0.253	0.204	0.207	0.198	
std	0.022	0.036	0.031	0.050	0.024	

Analysis of the results shows the redistribution of weights due to correlation between the attributes of the alternatives. There was an increase in the weight of 3rd criterion (weak correlation of attributes with the rest) and a weakening in the weight of 1st criterion (significant correlation of attributes with the rest). Comparison of the results with similar estimates by the Entropy and CRITIC methods (Table 2) does not reveal any definite pattern.

Since there are no criteria for the adequacy of various weight assessment methods, the effectiveness of the EWM-Corr modification is relative. The choice of a particular weighing method and its appropriateness is assessed by the presence of arguments “for and against”. This applies to all methods without exception. In our case, the argument is the redistribution of weights between correlated and uncorrelated attributes.

4.2. Assessment of the entropy of the criterion using the distribution function of the attribute of alternatives: EWM.df modification

After the decision matrix is normalized using the Max-Min transformation, the domains of the normalized values of attribute are consistent with each other. The priority of individual criteria due to different measurement scales is eliminated.

Information about the states of an object is determined by the distribution of values within the domain, not by the values themselves. Therefore, the calculation of entropy by values is not correct. A discrete random variable is determined by the pair (x_i, p_i) , and the values of the probabilities p_i are needed to calculate the entropy. To estimate the probabilities of possible states of alternatives for a fixed criterion, it is necessary to construct a discrete distribution using a statistical approach by performing the transition to frequency characteristics.

For the case of small samples, the distribution of a random variable can be determined based on estimates of the parameters of a hypothetical distribution of a random variable (for example, normal). The distribution assessment includes a statistical test for testing a hypothetical distribution law (for example, the Jarque-Bera test for checking normality (Jarque & Bera, 1987). Next, it is necessary to divide the domain of observed values into m intervals $[x_{ij}, x_{i+1,j}]$ and calculate the probability of the attribute falling into a certain interval. It does not matter that some of the intervals obtained by dividing the domain of values remain "unoccupied" by the actual values of the decision matrix. The frequency of the attribute values is taken into account through the distribution parameters. This does not exclude the existence of alternatives with such states. Potential probabilities of states are calculated by the expression:

$$p_{ij} = \int_{x_{ij}}^{x_{i+1,j}} f(x, \theta_1, \theta_2, \dots) dx, \quad (30)$$

where f is the density of the feature distribution, $\theta_1, \theta_2, \dots$ are the distribution parameters.

Below is a fragment of the calculation procedure for the case of a normal feature distribution law, performed in MatLab using built-in functions:

```
% define DM(m,n)
mu=mean(DM); sigma=std(DM);
mx=max(DM); mn=min(DM); h=(mx-mn)/(m-1);
for j=1:n
    [hJB, pJB] = jbttest(DM(:,j)); % Jarque-Bera test
    if hJB==0
        xN=mn(j)-h(j)/2 : h(j) : mx(j)+h(j)/2;
        xN1=xN(1:m);
        xN2=xN(2:m+1);
        pd = makedist('Normal',mu(j),sigma(j));
        y = cdf(pd,xN2)-cdf(pd,xN1); % cumulative distribution functions
    end
    pij(:,j)=y';
end
```

The calculations and illustration for the described algorithm are shown below in Figure 8 and Table 8.

When calculating the probability of possible states of alternatives, the mean value, standard deviation, the smallest and largest values of the attribute of alternatives are used. The results of evaluating the weight of the same criteria based on a different set of alternatives ($DM-1, DM-2, DM-3, DM-12$) are not so much different, as in the case of calculating the probabilities of possible states of alternatives using expression (1), in which a_{ij} is used to determine p_{ij} .

Specific character of objective methods for determining weights of criteria...

What to do in the absence of information about the distribution of features or rejection of the statistical test H_0 hypothesis of distribution?

It remains to apply an expert approach. The subjectivity of the assessment with a good expert is minimal. The expert's preferences are a reflection of the frequency of the feature, which is formed intuitively on the basis of experience. Expert scores are, in our opinion, better than statistics, which rarely have integrity and comprehensive coverage.




Figure 8. Probabilities of states of attributes of alternatives of the first criterion (normal distribution, Decision Matrix from Table 1).

Table 8. Estimation of the entropy and weights of the criterion using the EWM.df method.

Decision matrix	EWM.df					Rank (SAW)
	w_1	w_2	w_3	w_4	w_5	
DM-1	0.205	0.205	0.187	0.221	0.182	A2>A6>A8>...
DM-2	0.156	0.233	0.238	0.177	0.196	B5>B4>B2>...
DM-3	0.205	0.220	0.174	0.229	0.173	A2>A6>A8>...
DM-12	0.211	0.231	0.171	0.215	0.173	A2>A6>A8>...
mean	0.194	0.222	0.192	0.210	0.181	
std	0.025	0.013	0.031	0.023	0.011	
	Entropy (base)					
DM-1	0.146	0.228	0.211	0.223	0.191	A2>A4>A6>...
DM-2	0.107	0.287	0.240	0.134	0.232	B3>B7>B1>...
DM-3	0.144	0.216	0.200	0.249	0.191	A2>A4>B1>...
DM-12	0.159	0.280	0.165	0.220	0.176	B3>A4>A2>...
mean	0.139	0.253	0.204	0.207	0.198	
std	0.022	0.036	0.031	0.050	0.024	

4.3. Estimation of the entropy of the criterion using the relative disposition of alternatives: EWM.dsp modification

Let us arrange the values of the j -th attribute of alternatives in ascending order and calculate the relative disposition of the ordered set of alternatives for each j -th criterion using expression (13):

$$u_{ij} = \underset{i}{\text{sort}}(a_{ij}, 'ascend'), \tag{31}$$

$$d_j^{pq} = \frac{u_{qj} - u_{pj}}{u_j^{\max} - u_j^{\min}}, \forall p, q = 1, \dots, m \tag{32}$$

We get a discrete set $(m-1)$ of values:

$$d_j = \{ d_j^{1,2}, d_j^{2,3}, \dots, d_j^{(m-1),m} \} \tag{33}$$

d_j^{pq} are the normalized distances between adjacent alternatives (in an ordered list) for the j -th attribute.

To estimate the probabilities of the states of the j -th attribute for the EWM-method, we apply not the normalized values of the attributes p_{ij} , but a discrete set (33) consisting of $(m-1)$ the values of the relative dispositions of alternatives for each criterion.

Let's justify this decision:

1) set (33) characterizes the degree of order. If the distribution of the values of the j -th attribute is uniform:

$$d_j^{1,2} = d_j^{2,3} = \dots = d_j^{(m-1),m} = \frac{1}{m-1}, \tag{34}$$

then the entropy of the j -th criterion is $e_j = 1$, and its weight is 0. The order "in the line" is ideal when the distance between neighboring elements is equal. Any violation of the order leads to a decrease in entropy,

2) in accordance with expression (12), the p_{ij} values are the relative distances of the ordered list of alternatives of the j -th attribute to the alternative with the smallest value, i.e. tied to a reference point. In contrast to this, d_j^{pq} characterizes not the position, but the state, and do not depend on the position of the point on the scale [0; 1],

3) if the distribution of system states is uniform, then the values p_{ij} determine the probability that the random value X of the j -th attribute belongs to the interval $[a_j^{\min}; a_{ij}]$:

$$p_{ij} = P(a_j^{\min} < X < a_{ij}) = P(0 < X < p_{ij}) = \frac{a_{ij} - a_j^{\min}}{\sum_i (a_{ij} - a_j^{\min})} \tag{35}$$

p_{ij} depends on the position of the attribute on the scale [0; 1] and contradicts the concept of the probability of a state.

The d_j^{pq} values determine the probability that the random value X of the j -th attribute belongs to the interval $[u_{jp}; u_{jp}]$ for ordered set of alternatives:

Specific character of objective methods for determining weights of criteria...

$$d_j^{pq} = P(u_{pj} < X < u_{qj}) = \frac{u_{qj} - u_{pj}}{u_j^{\max} - u_j^{\min}} \quad (36)$$

and correspond to the concept of the probability of a state as the probability of belonging to a localized range of values,

4) as noted above, the relative disposition of the attribute of alternatives is invariant with respect to linear data transformation (according to the formula (13)) or retains almost all information about the system (minus two degrees of freedom),

5) each of the elements of the set (33) represents the intensity of the state

$$0 \leq d_j^{i-(i+1)} < 1, \quad \sum_{i=1}^{m-1} d_j^{i-(i+1)} = 1 \quad (37)$$

6) set (33) represents a differentiated series or the relative rate of change of values in individual sections,

7) unlike p_{ij} by the formula (12) the sets $\{d_j\}$ for different j are not subject to displacement relative to each other, and the size of the domains does not depend on the normalization of Sum,

8) set (33) can be considered as an analogue of the distribution histogram.

Thus, to calculate the entropy of each criterion, we use the quantities $\{d_j\}$:

$$e_j = -\frac{1}{\ln(m_j - 1)} \cdot \sum_{i=1}^{m-1} d_j^{i,(i+1)} \cdot \ln(d_j^{i,(i+1)}), \quad (\text{if } d_j^{pq}=0 \Rightarrow d_j^{pq} \cdot \ln(d_j^{pq})=0) \quad (38)$$

where m_j is dimension of the set $\{d_j^{pq}\}$ of different attribute (without 0) values by the j -th criterion (without zeros).

Next, we calculate the weights of the criteria using expression (6).

Notes:

- i. in the EWM.dsp method we used the concept that the probability of a state is proportional to the length of the localized gap. This concept is adequate for the even distribution of attributes. If the distribution is not uniform (or very different from uniform), then it is better to use the EWM.df approach described in section 4.2 above. In any case, EWM.dsp is better than estimates of the probability of states using expression (35) for the basic EWM-method.
- ii. the proposed approach does not require inversion of values for cost criteria,
- iii. if $e_j = 1$ for all j , then the weights should be taken equal,
- iv. if k alternatives have the same attributes, then $d_j^{pq} = 0$. $(k-1)$ values are excluded from the calculations and the entropy is calculated from the $(m-k)$ values.

The results of evaluating the entropy and weight of criteria by the proposed EWM.dsp method for matrices $DM-1$, $DM-2$, $DM-3$, $DM-12$ are presented in Table 9. The weights of the criteria are not as different as in the case of using the basic EWM-method (Table 2).

Figure 9 shows the summary probability diagrams of alternative states and entropy components ($-p \cdot \ln p$) for the decision matrix *DM-1*, obtained in various modifications of the EWM-method.

Table 9. Estimation of the entropy and weights of the criterion using the EWM.dsp method.

decision matrix	e_1	e_2	e_3	e_4	e_5	
DM-1	0.934	0.813	0.914	0.885	0.825	
DM-2	0.949	0.671	0.686	0.793	0.700	
DM-3	0.916	0.806	0.946	0.822	0.857	
DM-12	0.977	0.802	0.917	0.756	0.844	
mean	0.944	0.773	0.866	0.814	0.807	
std	0.026	0.068	0.121	0.054	0.072	
	w_1	w_2	w_3	w_4	w_5	Rank (SAW)
DM-1	0.105	0.298	0.137	0.182	0.278	A4>A2>A1>...
DM-2	0.043	0.274	0.261	0.172	0.250	B3>B7>B1>...
DM-3	0.128	0.297	0.082	0.273	0.220	A2>A4>B1>...
DM-12	0.033	0.281	0.118	0.347	0.221	A2>B3>A4>...
mean	0.077	0.288	0.150	0.244	0.242	
std	0.046	0.012	0.078	0.083	0.028	

Due to the equality of the attributes for some alternatives, in the calculations of the EWM.dsp method of the presented example, the number of states involved is less than $(m-1)$. The problem under study, defined by the *DM-1* matrix, is weakly sensitive to weight variation. Therefore, for all modifications of the EWM-method, the ranking for the studied example is the same.




Figure 9. Probabilities (p) of possible states of alternatives and entropy components ($-p \cdot \ln p$) of the Entropy method for the decision matrix *DM-1* for various modifications of the EWM-method. Two-axis technique.

Specific character of objective methods for determining weights of criteria...

5. Conclusion

The importance of the weights when aggregating the attributes of alternatives is such that alternatives with a priority on the attribute and a higher weight will receive priority in the performance indicator. Therefore, the quality of weight estimation for MCDM tasks is critical.

The conducted comprehensive analysis shows that the rationality of all objective methods for evaluating the criteria weights for MCDM tasks is questionable. Specific algorithms for objective methods for evaluating the weights of criteria still require further study.

The first decision that needs to be made and which will greatly affect the final results, is the choice between equal and different weights. Equal weighting is the preferred procedure in most applications and makes the weights estimation less subjective.

Differential weighing requires choosing the most appropriate approach for determining the weights. The solution should be supported by theoretical considerations that give meaning to each indicator or take into account its effect on synthesis in accordance with the structure of the problem.

It is important to compare weighting results for different methods. Since there are no criteria for the effectiveness of weighing methods, the discrepancy in the results of weighting requires serious analysis. The design and determination of weights can be interpreted in terms of value judgments, that methods based on the subjective opinions of individual experts are preferred.

It is obvious that the use of various objective (and other classes) methods and modifications lead in many cases to completely different values in the estimates of the weights of the criteria. There are also no criteria for the effectiveness of the methods. Therefore, further research, in our opinion, can be aimed at a constructive solution to the problem — building a decision support system (DSS) for weight estimation, including a wide range of methods, a knowledge base and an intelligent system for analyzing and synthesizing of results. In fact, it is an extended component of DSS for multi-criteria decision support systems.

All the algorithms described in this paper are implemented by the author in the MatLab system, posted in a file hosting service “File Exchange MathWorks” and are available for free use at the link (Math Works, 2021).

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflicts of interest.

References

- Ahn, B.S., & Park, K.S. (2008). Comparing methods for multiattribute decision making with ordinal weights. *Computers & Operations Research*, 35 (5), 1660–1670.
- Ali, Z., Mahmood, T., Ullah, K., & Khan, Q. (2021). Einstein Geometric Aggregation Operators using a Novel Complex Interval-valued Pythagorean Fuzzy Setting with Application in Green Supplier Chain Management. *Reports in Mechanical Engineering*, 2(1), 105-134.

Alosta, A., Elmansuri, O., & Badi, I. (2021). Resolving a location selection problem by means of an integrated AHP-RAFSI approach. *Reports in Mechanical Engineering*, 2(1), 135-142.

Barron, F.H. (1992). Selecting a best multiattribute alternative with partial information about attribute weights. *Acta Psychologica*, 80, 91-103.

Barron, F.H., & Barrett, B.E. (1996). Decision quality using ranked attribute weights. *Management Science*, 42, 1515-1523.

Barron, H., & Schmidt, C. (1988). Sensitivity analysis of additive multiattribute value models. *Operations Research*, 36(1), 122-127.

Bobko P., Roth, P.L., & Buster, M.A. (2007). The Usefulness of Unit Weights in Creating Composite Scores. A Literature Review, Application to Content Validity, and Meta-Analysis. *Organizational Research Methods*, 10(4), 689-709.

Chen, P. (2019). Effects of normalization on the entropy-based TOPSIS method. *Expert Systems with Applications*, 136, 33-41.

Đalić, I., Stević, Ž., Karamasa, C., & Puška, A. (2020) A Novel Integrated Fuzzy PIPRECIA – Interval Rough Saw Model: Green Supplier Selection. *Decision Making: Applications in Management and Engineering*, 3(1), 126-145.

Deng, M., Xu, W., & Yang, J.B. (2004). Estimating the attribute weights through evidential reasoning and mathematical programming. *International Journal of Information Technology & Decision Making*, 3, 419-428.

Diakoulaki, D., Mavrotas, G., & Papayannakis, L. (1995). Determining objective weights in multiple criteria problems: The CRITIC method. *Computers & Operations Research*, 22(7), 763-770.

Doyle, J.R., Green, R.H., & Bottomley, P.A. (1997). Judging relative importance: Direct rating and point allocation are not equivalent. *Organizational Behavior and Human Decision Processes*, 70, 55-72.

Ginevicius, R., & Podvezko, V. (2005). Objective and subjective approaches to determining the criterion weight in multicriteria models. *Proceedings of International Conference RelStat. Transport and Telecommunication*, 6(1), 133-137.

Goodwin, P., & Wright, G. (1998). *Decision Analysis for Management Judgment*. (2nd ed). Chichester: Wiley.

He, D., Xu, J., & Chen, X. (2016). Information-theoretic-entropy based weight aggregation method in multiple-attribute group decision-making. *Entropy*, 18(6), 171.

Hwang, C.L., & Yoon K. (1981). *Multiple Attributes Decision Making: Methods and Applications. A State-of-the-Art Survey*. Berlin: Springer-Verlag, Heidelberg, XI.

Jahan, A, & Edwards, K.L. (2015). A state-of-the-art survey on the influence of normalization techniques in ranking: Improving the materials selection process in engineering design. *Materials & Design*, 65, 335-342.

Jahan, A., Mustapha, F., Sapuan, S.M., Ismail, M. Y., Bahraminasab, M. (2012). A framework for weighting of criteria in ranking stage of material selection process. *The International Journal of Advanced Manufacturing Technology*, 58, 411-420.

Specific character of objective methods for determining weights of criteria...

Jarque, C. M., & Bera, A. K. (1987). A test for normality of observations and regression residuals. *International Statistical Review*, 55(2), 163–172.

Karande, P., Zavadskas, E., Chakraborty, S. (2016). A study on the ranking performance of some MCDM methods for industrial robot selection problems. *International Journal of Industrial Engineering Computations*, 7, 399–422.

Kersulienė, V., Zavadskas, E. K., & Turskis, Z. (2010). Selection of rational dispute resolution method by applying new step-wise weight assessment ratio analysis (SWARA). *Journal of Business Economics and Management*, 11(2), 243–258.

Kirkwood, C.W., & Corner, J.L. (1993). The effectiveness of partial information about attribute weights for ranking alternatives in multiattribute decision making. *Organizational Behavior and Human Decision Processes*, 54, 456–476.

Li, P., Qian, H., Wu, J., & Chen, J. (2013). Sensitivity analysis of TOPSIS method in water quality assessment: I. Sensitivity to the parameter weights. *Environmental Monitoring and Assessment*, 185, 2453–2461.

Li, X., Wang, K., Liu, L., Xin J., Yang, H., Gao, C. (2011). Application of the Entropy Weight and TOPSIS Method in Safety Evaluation of Coal Mines, *Procedia Engineering*, First International Symposium on Mine Safety Science and Engineering, 26, 2085–2091.

Liang, J., Shi, Z. D., Wierman, M.J. (2016). Information entropy, rough entropy and knowledge granulation in incomplete information systems. *International Journal of General Systems*, 35, 641–654.

Lotfi, F. H.; Fallahnejad, R. (2010). Imprecise Shannon's entropy and multi attribute decision making. *Entropy*, 12, 53–62.

Ma, J., Fan, Z. P., & Huang, L. H. (1999). A subjective and objective integrated approach to determine attribute weights. *European Journal of Operational Research*, 112, 397–404.

Mareschal, B. (1988). Weight stability intervals in multicriteria decision aid. *European Journal of Operational Research*, 33, 54–64.

Math Works. File Exchange. Weight of Criteria by Irik Mukhametzyanov (2021). <https://www.mathworks.com/matlabcentral/fileexchange/91720-weight-of-criteria> Accessed 06 May 2021.

Milosevic, T., Pamucar, D., Chatterjee, P. (2021). Model for selecting a route for the transport of hazardous materials using a fuzzy logic system. *Military Technical Courier*, 69(2), 355–390.

Mukhametzyanov, I. & Pamučar, D. (2018). Sensitivity analysis in MCDM problems: A statistical approach. *Decision Making: Applications in Management and Engineering*, 1(2), 51–80.

Mukhametzyanov, I. Z. (2018). IZ-norm method of normalization the multidimensional data, United States Copyright Office, Reg# TXu002125819/2018-09-28, <https://cocatalog.loc.gov/cgi-bin/Pwebrecon.cgi?DB=local&PAGE=First>

Mukhametzyanov, I. Z. (2019). A new method for the normalization of multidimensional data (IZ-Method) for MCDM problems, Proc. the 25st International Conference on Multiple Criteria Decision Making (MCDM 2019), Istanbul, Turkey (2019), 96–97.

Mukhametzyanov, I. Z. (2020). ReS-algorithm for converting normalized values of cost criteria into benefit criteria in MCDM tasks. *International Journal of Information Technology & Decision Making*, 19(5), 1389–1423.

Mukhametzyanov, I. Z. (2021). Elimination of the domains' displacement of the normalized values in MCDM tasks: the IZ-method. *International Journal of Information Technology & Decision Making*, [In Press].

Odu, G. O. (2019). Weighting Methods for Multi-Criteria Decision Making Technique. *Journal of Applied Sciences and Environmental Management*, 23(8), 1449–1457.

Pamučar D., Stević Ž., & Sremac S. (2018). A new model for determining weight coefficients of criteria in MCDM models: Full Consistency Method (FUCOM). *Symmetry*, 10(9), 393.

Pamucar, D., & Savin, L.M. (2020). Multiple-criteria model for optimal off-road vehicle selection for passenger transportation: BWM-COPRAS model. *Military Technical Courier*, 68(1), 28-64.

Pamucar, D., Ecer, F. (2020). Prioritizing the weights of the evaluation criteria under fuzziness: The fuzzy full consistency method – FUCOM-F. *Facta Universitatis, series: Mechanical Engineering*. 18(3), pp. 419 - 437.

Pekelman, D., & Sen, S.K. (1974). Mathematical programming models for the determination of attribute weights. *Management Science*, 20, 1217–1229.

Roberts, R., & Goodwin P. (2002). Weight approximations in multi-attribute decision models. *Journal of Multi-Criteria Decision Analysis*, 11, 291–303.

Saaty, T. L. (1980). *The Analytic Hierarchy Process*. New York: McGraw-Hill.

Shirland, L.E., Jesse, R.R., Thompson, R.L., & Iacovou, C.L. (2003). Determining attribute weights using mathematical programming. *Omega*, 31, 423–437.

Singh, A., Ghadai, R.K., Kalita, K., Chatterjee, P., Pamucar, D. (2020). EDM process parameter optimization for efficient machining of Inconel-718. *Facta Universitatis, series: Mechanical Engineering*. 18(3), pp. 473 - 490.

Solymosi, T., & Dombi, J. (1986). A method for determining the weights of criteria: The centralized weights. *European Journal of Operational Research*, 26, 35–41.

Stillwell, W. G., Seaver, D. A., & Edwards, W. (1981). A comparison of weight approximation techniques in multiattribute utility decision making. *Organizational Behavior and Human Performance*, 28, 62–77.

Takeda, E., Cogger, K. O., & Yu, P. L. (1987). Estimating criterion weights using eigenvectors: A comparative study. *European Journal of Operational Research*, 29, 360–369.

Triantaphyllou, E. (2000). *Multi-criteria Decision Making Methods: A Comparative Study* (Springer US).

Specific character of objective methods for determining weights of criteria...

Tzeng, G-H., & Huang, J-J. (2011). *Multiple Attribute Decision Making: Methods and Application*. Chapman and Hall/CRC.

Ustinovičius, L. (2001). Determining integrated weights of attributes. *Statyba*, 7(4), 321–326.

Vinogradova, I., Podvezko, V., & Zavadskas, E.K. (2018). The recalculation of the weights of criteria in MCDM methods using the bayes approach. *Symmetry*, 10(6), 205.

Von Winterfeldt, D., & Edwards, W. (1986). *Decision Analysis and Behavioral Research*, Cambridge: Cambridge University Press, MA.

Wang, Y.-M., & Luo, Y. (2010). Integration of correlations with standard deviations for determining attribute weights in multiple attribute decision making. *Math. Comput. Model.*, 51, 1–12.

Wolters, W. T. M., & Mareschal, B. (1995). Novel types of sensitivity analysis for additive MCDM methods, *European Journal of Operational Research*, 81(2), 281–290.

Wu, D., Wang, N., Yang, Z., Li, C., Yang, Y. (2018). Comprehensive Evaluation of Coal-Fired Power Units Using grey relational analysis and a hybrid entropy-based weighting method. *Entropy*, 20, 215.

Wu, J., Sun, J., Liang, L., & Zha, Y. (2011). Determination of weights for ultimate cross efficiency using Shannon entropy. *Expert Systems with Applications*, 38(5), 5162–5165.

Xu, X. (2004). A note on the subjective and objective integrated approach to determine attribute weights. *European Journal of Operational Research*, 156, 530–532.

Zeleny, M. (1982). *Multiple Criteria Decision Making*. McGraw-Hill, New York.

Zhao, H., Yao, L., Mei, G., Liu, T., Ning, Y. (2017). A Fuzzy comprehensive evaluation method based on AHP and entropy for landslide susceptibility map. *Entropy*, 19, 396.

Žižović, M., & Pamucar, D. (2019). New model for determining criteria weights: Level Based Weight Assessment (LBWA) model. *Decision Making: Applications in Management and Engineering*, 2(2), 126–137.

Žižović, M., Miljković, B., & Marinković, D. (2020). Objective methods for determining criteria weight coefficients: A modification of the CRITIC method. *Decision Making: Applications in Management and Engineering*, 3(2), 149–161.

Zolfani, S.H.; Yazdani, M.; Pamucar, D. & Zarate, P. (2020). A VIKOR and TOPSIS focused reanalysis of the MADM methods based on logarithmic normalization. *Facta universitatis series: Mechanical Engineering*, 18(3), 341–355.



© 2018 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).