

# MODIFICATION OF THE CRITIC METHOD USING FUZZY ROUGH NUMBERS

Dragan Pamucar<sup>1\*</sup>, Mališa Žižović<sup>2</sup> and Dragan Đuričić<sup>2</sup>

<sup>1</sup> Faculty of Organizational Sciences, University of Belgrade, Belgrade, Serbia

<sup>2</sup> University of Kragujevac, Kragujevac, Serbia

Received: 12 June 2022;

Accepted: 18 September 2022;

Available online: 16 October 2022.

*Original scientific paper*

**Abstract:** *This paper presents a new approach in the modification of the CRiteria Importance Through Intercriteria Correlation (CRITIC) method using fuzzy rough numbers. In the modified CRITIC method (CRITIC-M), the normalization procedure of the home matrix elements was improved and the aggregation function for information processing in the normalized home matrix was improved. By introducing a new way of normalization, smaller deviations between normalized elements are obtained, which affects smaller values of standard deviation. Thus, the relationships between the data in the initial decision matrix are presented in a more objective way. The introduction of a new way of aggregating the values of weights in the CRITIC-M method enables a more comprehensive view of information in the initial decision matrix, which leads to obtaining more objective values of weights. A new concept of fuzzy rough numbers was used to address uncertainties in the CRITIC-M methodology.*

**Key words:** *MCDM, fuzzy sets, rough sets, fuzzy rough numbers, CRITIC-M.*

## 1. Introduction

Determining criterion weights is one of the key problems that arises in multi-criteria optimization models. In order to develop effective methods for determining the weight of the criteria, researchers around the world in recent years in the literature pay considerable attention to this problem. Most authors suggest dividing the model for determining the weights of criteria into subjective and objective (Zhu et al., 2015).

Subjective approaches reflect the subjective opinion and intuition of the decision maker. In this approach, the weight of the criteria are determined based on the preferences of the decision maker. Traditional methods of determining weights of criteria include tradeoff method (Keeney & Raiffa, 1976), proportional (ratio) method, Swing method (Weber et al., 1988) and Conjoint method (Green & Srinivasan,

\* Corresponding author.

E-mail addresses: [dpamucar@gmail.com](mailto:dpamucar@gmail.com) (D. Pamucar), [zizovic@gmail.com](mailto:zizovic@gmail.com) (M. Žižović), [dragandjurcic@gmail.com](mailto:dragandjurcic@gmail.com) (D. Đuričić)

1990), Analytic Hierarchy Process (AHP) model (Saaty, 1980), SMART (the Simple Multi Attribute Rating Technique) method (Keeney & Raiffa, 1976), MACBETH (Measuring Attractiveness by Categorical Based Evaluation Technique) method (Bana e Costa & Vansnick, 1994), Direct point allocation method (Poyhonen & Hamalainen, 2001), Ratio or direct significance weighting method (Weber & Borchering, 1993), Resistance to change method (Rogers & Bruen, 1998), WLS (Weighted Lest Square) method (Graham, 1987) and FPP (the Fuzzy Preference Programming method) method (Mikhailov, 2000). Recent subjective methods include multipurpose linear programming (Costa & Climaco, 1999), linear programming (Mousseau et al., 2000), SWARA (Step-wise Weight Assessment Ratio Analysis) method (Valipour et al., 2017), BWM (Best Worst Method) (Rezaei, 2015) and FUCOM (FULL Consistency Method) (Pamučar et al., 20018).

Among the most known objective methods are the following: Entropy method (Shannon & Weaver, 1947), CRITIC method (CRiteria significance Through Intercriteria Correlation) (Srđević et al., 2003) and FANMA method whose name was derived from the names of the authors of the method (Žižović et al., 2020).

The CRITIC method is one of the most well-known and most frequently used objective methods. The CRITIC method belongs to the group of correlation methods, which uses standard deviations of the standardized criterion values of variants to determine the contrast of criteria, as well as the correlation coefficients of all pairs of columns. In this study, certain limitations were identified when applying the classical CRITIC method and a modification of the CRITIC method (CRITIC-M) in a fuzzy rough environment was proposed.

The rest of the work is organized as follows. The following section shows the preliminary settings for fuzzy rough numbers. Section 3 presents the mathematical foundations of the classical CRITIC method. While section 4 shows a modification of the CTIRIC method in a fuzzy rough environment. The fifth section of the paper presents the application of the fuzzy rough CRITIC-M method through an example from the literature. Concluding remarks and directions for future research are given in Section 6.

## 2. Preliminaries on fuzzy rough numbers

In the fuzzy rough concept, fuzzy theory was used to represent uncertainty in information, while rough theory was used to create flexible boundary intervals of fuzzy numbers. The use of hybrid fuzzy rough numbers eliminates the limitation of classic fuzzy type 2 numbers that have a predefined imprint of uncertainty.

We assume that  $U$  universe contains all of the objects and let  $Y$  be an arbitrary object from  $U$ . We assume there is a set of  $k$  classes which represent the preferences of the DM,  $G^* = (A_1, A_2, \dots, A_k)$ , with the condition that they belong to a series which satisfies the condition  $A_1 < A_2 < \dots < A_k$ . All objects are defined in the universe and connected with the preferences of the DM. Each element  $A_i$  ( $1 \leq i \leq k$ ) represents a fuzzy number that is defined as  $A_q = (a_{1q}, a_{2q}, a_{3q})$ . Since element  $A_i$  from the class of objects  $G^*$  is represented as fuzzy number  $A_q = (a_{1q}, a_{2q}, a_{3q})$ , for each value  $a_{1q}$ ,  $a_{2q}$  and  $a_{3q}$  we obtain one class of objects that is represented in the interval  $I(a_1)_q = \{I(a_1)_{lq}, I(a_1)_{uq}\}$ ,  $I(a_2)_q = \{I(a_2)_{lq}, I(a_2)_{uq}\}$  and  $I(a_3)_q = \{I(a_3)_{lq}, I(a_3)_{uq}\}$  where the condition is fulfilled that  $I(a_j)_{lq} \leq I(a_j)_{uq}$  ( $j = 1, 2, 3; 1 \leq q \leq k$ ), as well as the condition

$I(a_1)_q, I(a_2)_q, I(a_3)_q \in G^*$ . Then  $I(a_j)_{lq}$  and  $I(a_j)_{uq}$  ( $j=1,2,3; 1 \leq q \leq k$ ) respectively represent the lower and upper border of the intervals of the  $q$ -th class of objects. If both limits of the classes of objects (upper and lower limits) respectively are compared so that  $I^*(a_j)_{l1} < I^*(a_j)_{l2} < \dots < I^*(a_j)_{ls}; I^*(a_j)_{u1} < I^*(a_j)_{u2} < \dots < I^*(a_j)_{um}$  ( $j=1,2,3; 1 \leq s, m \leq k$ ), then for any of the classes of objects  $I^*(a_j)_{lq} \in G^*$  and  $I^*(a_j)_{uq} \in G^*$  ( $j=1,2,3; 1 \leq q \leq k$ ) we can define the lower approximation  $I^*(a_j)_{lq}$  using the following equations

$$\underline{Apr}(I^*(a_1)_{lq}) = \cup \{Y \in U / G^*(Y) \leq I^*(a_1)_{lq}\}; (1 \leq q \leq k) \tag{1}$$

$$\underline{Apr}(I^*(a_2)_{lq}) = \cup \{Y \in U / G^*(Y) \leq I^*(a_2)_{lq}\}; (1 \leq q \leq k) \tag{2}$$

$$\underline{Apr}(I^*(a_3)_{lq}) = \cup \{Y \in U / G^*(Y) \leq I^*(a_3)_{lq}\}; (1 \leq q \leq k) \tag{3}$$

And the upper approximation of  $I^*(a_j)_{uq}$  using the following equations

$$\overline{Apr}(I^*(a_1)_{uq}) = \cup \{Y \in U / G^*(Y) \geq I^*(a_1)_{uq}\}; (1 \leq q \leq k) \tag{4}$$

$$\overline{Apr}(I^*(a_2)_{uq}) = \cup \{Y \in U / G^*(Y) \geq I^*(a_2)_{uq}\}; (1 \leq q \leq k) \tag{5}$$

$$\overline{Apr}(I^*(a_3)_{uq}) = \cup \{Y \in U / G^*(Y) \geq I^*(a_3)_{uq}\}; (1 \leq q \leq k) \tag{6}$$

Both classes of objects (object classes  $I^*(a_j)_{lq}$  and  $I^*(a_j)_{uq}$ ) are defined by their lower limits  $\underline{Lim}(I^*(a_j)_{lq}); j=1,2,3$ , and upper limits  $\overline{Lim}(I^*(a_j)_{uq}); j=1,2,3$ . The lower limits are defined by the following equations

$$\underline{Lim}(I^*(a_1)_{lq}) = \frac{1}{M_{L(a_1)}} \sum G^*(Y) | Y \in \underline{Apr}(I^*(a_1)_{lq}); (1 \leq q \leq k) \tag{7}$$

$$\underline{Lim}(I^*(a_2)_{lq}) = \frac{1}{M_{L(a_2)}} \sum G^*(Y) | Y \in \underline{Apr}(I^*(a_2)_{lq}); (1 \leq q \leq k) \tag{8}$$

$$\underline{Lim}(I^*(a_3)_{lq}) = \frac{1}{M_{L(a_3)}} \sum G^*(Y) | Y \in \underline{Apr}(I^*(a_3)_{lq}); (1 \leq q \leq k) \tag{9}$$

where  $M_{L(a_1)}, M_{L(a_2)}$  and  $M_{L(a_3)}$  respectively represent the number of objects included in the lower approximation of the classes of objects  $I^*(a_1)_{lq}, I^*(a_2)_{lq}$  and  $I^*(a_3)_{lq}$ . The upper limits  $\overline{Lim}(I^*(a_j)_{uq}); j=1,2,3$  are defined by equations (10)-(12)

$$\overline{Lim}(I^*(a_1)_{uq}) = \frac{1}{M_{U(a_1)}} \sum G^*(Y) | Y \in \overline{Apr}(I^*(a_1)_{uq}); (1 \leq q \leq k) \tag{10}$$

$$\overline{Lim}(I^*(a_2)_{uq}) = \frac{1}{M_{U(a_2)}} \sum G^*(Y) | Y \in \overline{Apr}(I^*(a_2)_{uq}); (1 \leq q \leq k) \tag{11}$$

$$\overline{Lim}(I^*(a_3)_{uq}) = \frac{1}{M_{U(a_3)}} \sum G^*(Y) | Y \in \overline{Apr}(I^*(a_3)_{uq}); (1 \leq q \leq k) \tag{12}$$

where  $M_{U(a_1)}, M_{U(a_2)}$  and  $M_{U(a_3)}$  respectively represent the number of objects that are contained in the upper approximation of the classes of objects  $I^*(a_1)_{uq}, I^*(a_2)_{uq}$  and  $I^*(a_3)_{uq}$ .

As we see, each class of objects  $I(a_1)_q, I(a_2)_q$  and  $I(a_3)_q$  is defined by means of its own lower and upper limits, which make up the interval fuzzy-rough number  $\overline{A}$  Figure 1, defined as

Modification of the CRITIC method using fuzzy rough numbers

$$\overline{\overline{A}} = [A_q^L, A_q^U] = \left[ \left( \overline{\text{Lim}}(I^*(a_1)_{uq}), \underline{\text{Lim}}(I^*(a_2)_{lq}), \overline{\text{Lim}}(I^*(a_2)_{uq}), \underline{\text{Lim}}(I^*(a_3)_{lq}); w_1(A_q^L) \right), \left( \underline{\text{Lim}}(I^*(a_1)_{lq}), \overline{\text{Lim}}(I^*(a_2)_{lq}), \overline{\text{Lim}}(I^*(a_2)_{uq}), \underline{\text{Lim}}(I^*(a_3)_{uq}); w_2(A_q^U) \right) \right] \quad (13)$$

where  $A_q^L$  and  $A_q^U$  respectively represent the upper and lower trapezoidal fuzzy-rough number which meets the condition that  $A_q^L \subset A_q^U$ , while  $w_1(A_q^L)$  and  $w_2(A_q^U)$  respectively represent the maximum values of interval fuzzy-rough number  $\overline{\overline{A}}$ .




Figure 1. Interval fuzzy-rough number  $\overline{\overline{A}}$

From Figure 1 we observe that for interval-valued fuzzy-rough number  $\overline{\overline{A}}$  it is valid that  $w_1(A_q^L) = w_2(A_q^U) = 1$ . On this basis we can write equation (13) in the following form:

$$\overline{\overline{A}} = [A_q^L, A_q^U] = \left[ (a_{1q}^L, a_{1q}^U), (a_{2q}^L, a_{2q}^U), (a_{3q}^L, a_{3q}^U) \right] \quad (14)$$

where  $a_{jq}^L = \underline{\text{Lim}}(I^*(a_j)_{lq})$  and  $a_{jq}^U = \overline{\text{Lim}}(I^*(a_j)_{uq})$ ; ( $j = 1, 2, 3$ ;  $1 \leq q \leq k$ ).

If there is consensus among the decision makers on the assignment of specific values from the linguistic fuzzy scale then  $a_{1q}^L = a_{1q}^U$ ,  $a_{2q}^L = a_{2q}^U$  and  $a_{3q}^L = a_{3q}^U$ . Then interval-valued fuzzy-rough number  $\overline{\overline{A}}$  becomes fuzzy number  $A$  type-1.

### 3. CRITIC method

The CRITIC method (CRiteria Importance Through Intercriteria Correlation) (Žižović et al., 2020) is a correlation method. Standard deviations of ranked criteria values of options in columns, as well as correlation coefficients of all paired columns are used to determine criteria contrasts.

Step 1: Starting from an initial decision matrix,  $X = [\xi_{ij}]_{m \times n}$ , we normalize the element of the initial decision matrix and form the normalized matrix  $X = [\hat{\xi}_{ij}]_{m \times n}$ .

$$X = \begin{matrix} & C_1 & C_2 & \dots & C_n \\ \begin{matrix} A_1 \\ A_2 \\ \vdots \\ A_m \end{matrix} & \begin{bmatrix} \xi_{11} & \xi_{12} & \dots & \xi_{1n} \\ \xi_{21} & \xi_{22} & \dots & \xi_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ \xi_{m1} & \xi_{m2} & \dots & \xi_{mn} \end{bmatrix} \end{matrix} \quad (15)$$

The normalization of matrix elements  $X = [\xi_{ij}]_{m \times n}$  is done by applying (16) and (17):

a) for maximizing criteria:

$$\xi_{ij} = \frac{\xi_{ij} - \xi_j^{\min}}{\xi_j^{\max} - \xi_j^{\min}}, \quad i = 1, 2, \dots, n; \quad j = 1, 2, \dots, m; \quad (16)$$

b) for minimizing criteria:

$$\xi_{ij} = \frac{\xi_j^{\max} - \xi_{ij}}{\xi_j^{\max} - \xi_j^{\min}}, \quad i = 1, 2, \dots, n; \quad j = 1, 2, \dots, m; \quad (17)$$

where  $\xi_j^{\max} = \max\{\xi_{1j}, \xi_{2j}, \dots, \xi_{mj}\}$ ;  $\xi_j^{\min} = \min\{\xi_{1j}, \xi_{2j}, \dots, \xi_{mj}\}$ .

Upon normalizing criteria of the initial decision matrix, all elements  $\xi_{ij}$  are reduced to interval values [0, 1], so it can be said that all criteria have the same metrics.

*Step 2:* For criterion  $C_j$  ( $j = 1, 2, \dots, n$ ) we define the standard deviation  $\sigma_j$ , that represents the measure of deviation of values of alternatives for the given criterion of average value. Standard deviation of a given criterion is the measure considered in the further process of defining criteria weight coefficients.

*Step 3:* From the normalized matrix  $X = [\xi_{ij}]_{m \times n}$  we separate the vector  $\xi_j = (\xi_{1j}, \xi_{2j}, \dots, \xi_{mj})$  that contains the values of alternatives  $A_i$  ( $i = 1, 2, \dots, m$ ) for the given criterion  $C_j$  ( $j = 1, 2, \dots, n$ ). After forming the vector  $\xi_j = (\xi_{1j}, \xi_{2j}, \dots, \xi_{mj})$ , we construct the matrix  $L = [l_{jk}]_{n \times n}$ , that contains coefficients of linear correlation of vectors  $\xi_j$  and  $\xi_k$ .

The quantity of data  $W_j$  contained within criterion  $j$  is determined by combining previously listed measures  $\sigma_j$  and  $l_{jk}$  as follows:

$$W_j = \sigma_j \cdot \varphi_j = \sigma_j \sum_{k=1}^n (1 - l_{kj}) \quad (18)$$

Based on the previous analysis we can conclude da a higher value  $W_j$  means a larger quantity of data received from a given criterion, which in turn increases the relative significance of the given criterion for the given decision process.

*Step 4:* Objective weights of criteria are reached by normalizing measures  $W_j$ :

$$w_j = \frac{W_j}{\sum_{k=1}^m W_k} \quad (19)$$

#### 4. Fuzzy rough CRITIC method

The modification of the CRITIC method presented in this section is based on three starting points: 1) Modification of the initial decision matrix data normalization method, 2) Modification of the expression for determining the final values of criterion weights, and 3) Extension of the modified CRITIC method using fuzzy rough numbers. In the following part, the modified fuzzy rough CRITIC method algorithm is presented and testing is performed on an example from the literature.

*Step 1. Construct the basic fuzzy rough decision matrix ( $\mathfrak{Z}$ ).* We will assume that the evaluation of alternatives was performed by  $e$  experts using the fuzzy scale. Also, we will assume that expert preferences are presented in the home matrix  $\mathfrak{Z}^b = \left[ \mathcal{G}_{ij}^b \right]_{m \times n}$

where  $1 \leq b \leq e$ ;  $i=1, \dots, m$ ;  $j=1, \dots, n$ ; and  $\mathcal{G}_{ij}^b = \left( \mathcal{G}_{ij}^{b(l)}, \mathcal{G}_{ij}^{b(m)}, \mathcal{G}_{ij}^{b(u)} \right)$  represent linguistic variables from the fuzzy scale used by expert  $e$ . For each element  $\theta_{ij}^{e(l)}$ ,  $\theta_{ij}^{e(m)}$  and  $\theta_{ij}^{e(u)}$

from  $\mathfrak{Z}^b = \left[ \mathcal{G}_{ij}^b \right]_{m \times n}$  we form matrices of the aggregated sequences of experts

$\mathfrak{Z}^{b(l)} = \left[ \mathcal{G}_{ij}^{b(l)} \right]_{m \times n}$ ,  $\mathfrak{Z}^{b(m)} = \left[ \mathcal{G}_{ij}^{b(m)} \right]_{m \times n}$  and  $\mathfrak{Z}^{b(u)} = \left[ \mathcal{G}_{ij}^{b(u)} \right]_{m \times n}$ . Using expressions (1)-(12)

sequence  $\mathcal{G}_{ij}^{e(l)}$ ,  $\mathcal{G}_{ij}^{e(m)}$  and  $\mathcal{G}_{ij}^{e(u)}$  are transformed into fuzzy rough number

$\bar{\mathcal{G}}_{ij}^b = \left( \left[ \bar{\mathcal{G}}_{ij}^{b(l)-}, \bar{\mathcal{G}}_{ij}^{b(l)+} \right], \left[ \bar{\mathcal{G}}_{ij}^{b(m)-}, \bar{\mathcal{G}}_{ij}^{b(m)+} \right], \left[ \bar{\mathcal{G}}_{ij}^{b(u)-}, \bar{\mathcal{G}}_{ij}^{b(u)+} \right] \right)$ ;  $1 \leq b \leq e$ . For fusion fuzzy rough values

$\bar{\mathcal{G}}_{ij}^b$  ( $1 \leq b \leq e$ ) the fuzzy rough weighted geometric Bonferroni function was used. This

is how the aggregated fuzzy rough matrix  $\bar{\mathfrak{Z}} = \left[ \bar{\mathcal{G}}_{ij} \right]_{m \times n}$  is defined.

*Step 2.* The elements of the matrix  $\bar{\mathfrak{Z}} = \left[ \bar{\mathcal{G}}_{ij} \right]_{m \times n}$  are normalized as follows:

$$\bar{\xi}_{ij} = \begin{cases} \left[ \left( \frac{\bar{\mathcal{G}}_{ij}^{(l)-}}{\mathcal{G}_{j+}^U}, \frac{\bar{\mathcal{G}}_{ij}^{(l)+}}{\mathcal{G}_{j+}^U} \right), \left( \frac{\bar{\mathcal{G}}_{ij}^{(m)-}}{\mathcal{G}_{j+}^U}, \frac{\bar{\mathcal{G}}_{ij}^{(m)+}}{\mathcal{G}_{j+}^U} \right), \left( \frac{\bar{\mathcal{G}}_{ij}^{(u)-}}{\mathcal{G}_{j+}^U}, \frac{\bar{\mathcal{G}}_{ij}^{(u)+}}{\mathcal{G}_{j+}^U} \right) \right]; & \text{if } j \in B, \\ \left[ \left( \frac{\mathcal{G}_{j-}^U}{\bar{\mathcal{G}}_{ij}^{(u)+}}, \frac{\mathcal{G}_{j-}^U}{\bar{\mathcal{G}}_{ij}^{(u)-}} \right), \left( \frac{\mathcal{G}_{j-}^U}{\bar{\mathcal{G}}_{ij}^{(m)+}}, \frac{\mathcal{G}_{j-}^U}{\bar{\mathcal{G}}_{ij}^{(m)-}} \right), \left( \frac{\mathcal{G}_{j-}^U}{\bar{\mathcal{G}}_{ij}^{(l)+}}, \frac{\mathcal{G}_{j-}^U}{\bar{\mathcal{G}}_{ij}^{(l)-}} \right) \right]; & \text{if } j \in C \end{cases} \quad (20)$$

where  $\mathcal{G}_{i+}^U = \max_{1 \leq i \leq m} (\mathcal{G}_{ij}^{(u)+})$  and  $\mathcal{G}_{i-}^L = \min_{1 \leq i \leq m} (\mathcal{G}_{ij}^{(l)-})$ .

*Step 3:* Construct a matrix of linear correlations. For each criterion  $C_j$  from the normalized matrix  $\bar{\mathfrak{Z}}^N = \left[ \bar{\xi}_{ij} \right]_{m \times n}$ , the vector  $\bar{\xi}_j = \left( \bar{\xi}_{1j}, \bar{\xi}_{2j}, \dots, \bar{\xi}_{mj} \right)$  is defined, and linear

correlations of the vectors  $\bar{\xi}_j$  and  $\bar{\xi}_k$  are calculated. By summing the linear correlations by criteria, we obtain the measure of the conflict of criteria

$\varphi_j = \sum_{k=1}^n (1 - l_{jk})$ . The amount of information  $W_j$  contained in criterion  $j$  is determined

by applying expression (21):

$$W_j = \sigma_j \sum_{k=1}^n (1 - l_{kj}) \quad (21)$$

*Korak 4:* Determination of weight coefficients of criteria. Objective weights of criteria are obtained by applying expression (22):

$$w_j = \frac{\frac{\bar{\xi}_j}{1 - \bar{\xi}_j} \cdot W_j}{\sum_{j=1}^n \left( \frac{\bar{\xi}_j}{1 - \bar{\xi}_j} \cdot W_j \right)} \tag{22}$$

Example:

We will assume that the multi-criteria model considers the evaluation of three alternatives under five criteria. We will also assume five experts evaluated the alternatives using the fuzzy scale presented in Table 1.

Table 1. Fuzzy scale

| Linguistic terms     | Membership function |
|----------------------|---------------------|
| Absolutely low (AL)  | (1, 1.5, 2.5)       |
| Very low (VL)        | (1.5, 2.5, 3.5)     |
| Low (L)              | (2.5, 3.5, 4.5)     |
| Medium low (ML)      | (3.5, 4.5, 5.5)     |
| Equal (E)            | (4.5, 5.5, 6.5)     |
| Medium high (MH)     | (5.5, 6.5, 7.5)     |
| High (H)             | (6.5, 7.5, 8.5)     |
| Extremely high (EH)  | (7.5, 8.5, 9.5)     |
| Absolutely high (AH) | (8.5, 9, 10)        |

Experts' assessments of alternatives are presented in Table 2.

Table 2. Expert evaluation of alternatives

|    | A1             | A2           | A3             |
|----|----------------|--------------|----------------|
| C1 | EH,EH,EH,AH,AH | H,EH,H,MH,H  | VL,L,L,L,L     |
| C2 | AH,AH,AH,AH,EH | E,ML,ML,ML,E | AH,AH,AH,H,AH  |
| C3 | EH,AH,AH,EH,AH | H,EH,H,EH,H  | EH,H,H,EH,AH   |
| C4 | EH,AH,AH,H,AH  | H,H,H,H,EH   | MH,MH,E,MH,E   |
| C5 | VL,VL,AL,VL,VL | E,E,ML,ML,ML | AL,AL,VL,VL,AL |

By applying expressions (1) - (12) the expert estimates were transformed into fuzzy rough values, Table 3.

Table 3. Fuzzy rough home matrix

| Crit. | A1                                    | A2                                    |
|-------|---------------------------------------|---------------------------------------|
| C1    | [[7.56,8.28],[8.53,8.89],[9.53,9.89]] | [[5.96,6.75],[6.97,7.76],[7.97,8.76]] |
| C2    | [[7.97,8.50],[8.74,9.00],[9.74,10.0]] | [[3.56,4.13],[4.56,5.13],[5.56,6.14]] |
| C3    | [[7.70,8.43],[8.60,8.97],[9.60,9.97]] | [[6.56,7.14],[7.56,8.14],[8.56,9.14]] |
| C4    | [[7.16,8.43],[8.07,8.97],[9.07,9.97]] | [[6.50,6.81],[7.5,7.81],[8.50,8.810]] |
| C5    | [[1.22,1.50],[1.93,2.50],[2.95,3.50]] | [[3.56,4.13],[4.56,5.13],[5.56,6.14]] |
| Crit. | A3                                    |                                       |
| C1    | [[1.93,2.44],[2.95,3.44],[3.96,4.45]] |                                       |
| C2    | [[7.41,8.39],[8.19,8.92],[9.20,9.92]] |                                       |
| C3    | [[6.70,7.64],[7.70,8.49],[8.70,9.49]] |                                       |
| C4    | [[4.70,5.33],[5.70,6.33],[6.70,7.33]] |                                       |
| C5    | [[1.03,1.31],[1.55,2.11],[2.56,3.12]] |                                       |

Using the expression (20), the elements from Table 3 were normalized. Then, using the expressions (21) and (22), the matrices of linear correlations of fuzzy rough elements were defined and the final values of the weighting coefficients were determined as follows:

$$w_1 = 0.153;$$

$$w_2 = 0.380;$$

$$w_3 = 0.189;$$

$$w_4 = 0.118;$$

$$w_5 = 0.160.$$

## 5. Conclusion

This research presents a modification of the CRITIC method using fuzzy rough numbers. Fuzzy rough numbers are applied because part of the uncertainty and subjectivity are neglected in the classic fuzzy and rough models. Given the well-known performance of fuzzy sets in representing uncertainties and confirmed advantages of rough numbers in subjectivity manipulation, a modification of the CRITIC method based on information processing using hybrid fuzzy rough numbers is proposed. Also, the application of the fuzzy rough CRITIC method is shown in an example that considers the evaluation of three alternatives under five criteria.

**Author Contributions:** Conceptualization, D.P. and M.Ž.; methodology, D.P., M.Ž. and D.Đ.; software, D.P., M.Ž. and D.Đ.; validation, D.P., M.Ž. and D.Đ.; formal analysis, D.P.; investigation, D.P., M.Ž. and D.Đ.; writing—original draft preparation, D.P. and M.Ž.; writing—review and editing, D.P., M.Ž. and D.Đ.; visualization, D.P.; supervision, D.P. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Conflicts of Interest:** The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.



## References

- Bana e Costa, C., & Vansnick, J.C. (1994). MACBETH: An interactive path towards the construction of cardinal value functions, *International Transactions in Operational Research*, 1(4), 489–500.
- Costa, J.P., & Climaco, J.C. (1999). Relating reference points and weights in MOLP. *Journal of Multi-Criteria Decision Analysis*, 8, 281-290.
- Graham, A. (1987). *Nonnegative matrices and applicable topics in linear algebra*, Ellis Horwood, Chichester, UK.
- Green, P.E., & Srinivasan, V. (1990). Conjoint analysis in marketing: New developments with implications for research and practice. *Journal of Marketing*, 54, 3-19.
- Keeney, R.L., & Raiffa, H. (1976). *Decisions with multiple objectives*. Wiley, New York.
- Mikhailov, L. (2000). A fuzzy programming method for deriving priorities in the analytic hierarchy process, *Journal of Operational Research Society*, 51, 341–349.
- Mousseau, V., Slowinski, R., & Zielniewicz, P. (2000). A user-oriented implementation of the ELECTRE-TRI method integrating preference elicitation support. *Computers and Operations Research*, 27, 757-777.
- Pamučar D, Stević, Ž., & Sremac, S. (2018). A New Model for Determining Weight Coefficients of Criteria in MCDM Models: Full Consistency Method (FUCOM). *Symmetry*, 10(9), 393.
- Poyhonen, M., & Hamalainen, R. (2001). On the convergence of multiattribute weighting methods, *European Journal of Operational Research*, 129, 569–585.
- Rezaei, J. (2015). Best-worst multi-criteria decision-making method. *Omega*, 53, 49-57.
- Rogers, M., & Bruen, M. (1998). A new system for weighting environmental criteria for use within ELECTRE III, *European Journal of Operational Research*, 107(3), 552–563.
- Saaty, T.L. (1980). *Analytic hierarchy process*. McGraw-Hill, New York.
- Shannon, C.E., & Weaver, W. (1947). *The mathematical theory of communication*. Urbana: The University of Illinois Press.
- Srđević, B., Medeiros, Y.D.P., Faria, A.S., & Schaer M. (2003). Objektivno vrednovanje kriterijuma performanse sistema akumulacija. *Vodoprivreda*, 35, 163-176. (Only in Serbian)
- Valipour, A., Yahaya, N., Md Noor, N., Antuchevičienė, J., & Tamošaitienė, J. (2017). Hybrid SWARA-COPRAS method for risk assessment in deep foundation excavation project: An Iranian case study. *Journal of Civil Engineering and Management*, 23(4), 524-532.
- Weber, M., & Borcherding, K. (1993). Behavioral influences on weight judgments in multiattribute decision making, *European Journal of Operational Research*, 67, 1–12.

#### Modification of the CRITIC method using fuzzy rough numbers

Weber, M., Eisenfuhr, F., & von Winterfeldt, D. (1988). The Effects of Splitting Attributes on Weights in Multiattribute Utility Measurement. *Management Science*, 34(2), 431-445.

Zhu, G.N., Hu, J., Qi, J., Gu, C.C., & Peng, J.H. (2015). An integrated AHP and VIKOR for design concept evaluation based on rough number, *Advanced Engineering Informatics*, 29, 408–418.

Žižović, M., Miljković, B., & Marinković, D. (2020). Objective methods for determining criteria weight coefficients: A modification of the CRITIC method. *Decision Making: Applications in Management and Engineering*, 3(2), 149-161.



© 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).