



CORRESPONDING AUTHOR

Marco Tecilla,
marco.tecilla@unimi.it

TeV3 outbreak characterization in captive *Testudo spp.*

M. Tecilla¹, P. Roccabianca¹, G. Grilli¹, F. Forlani¹, F.C. Origgi²

¹Department of Veterinary Science and Public Health, Università degli Studi di Milano, Via Celoria 10, 20133 Milan, Italy

²Center for Fish and Wildlife health, Vetsuisse-Faculty of Bern, Laenggassstrasse 122, 3012 Bern, Switzerland

ABSTRACT

Italian Tortoises species are considered either endangered or near threatened according to International Union for Conservation of Nature. When pet tortoises are abandoned or found injured or seized following illegal detention, they are sent to wildlife rehabilitation centers. From 2008, the *Testudo spp.* population housed in the WWF Vanzago's oasis exhibited clinical signs compatible with *Testudinid herpesvirus 3* (TeV3) infection. By the end of 2012 all *Testudo* had died. The presence of TeV3 was investigated by molecular biology and pathology. All the tortoises housed in Vanzago resulted ELISA positive for the presence of anti-TeV3 antibodies except one *T. hermanni*. Of these, 12 animals died and were all necropsied. Lesion frequency distribution was evaluated by histology. PCR was positive in 8/12 tortoises. To better complement the epidemiological evaluation of the virus in northern Italy, 20 retrospective cases were selected from the archive of the University of Milan. Of these, 5 were TeV3 PCR positive. Lesions closely resembled those of the Vanzago's population. These results are consistent with a high prevalence of TeV3 in northern Italy. The finding of intranuclear inclusion bodies demonstrated to be specific but not sensitive. TeV3 diagnostic pathological lesions have been reported to vary according with host immune response and by the viral replicative status. Molecular techniques were often necessary to confirm the infection. According to the literature and to our findings, *T. hermanni spp.* seems the species with higher mortality and lower antibody concentrations when infected with TeV3.

REFERENCES

- F. C. Origgi. 2012. *Journal of Herpetological Medicine and Surgery* 22 (1-2): 52-54; E. R. Jacobson., K. H. Berry, F. C. Origgi, J. F. Jr Wellehan, A. L. Childress, J. Braun, M. Schrenzel, J. Ye, B. Rideout. 2012. *Journal of Wildlife Diseases* 48(3): 747-57; F. C. Origgi, C. H. Romero, D. C. Bloom, P. A. Kelin, J. M. Gaskin, S. J. Tucker, E. R. Jacobson. 2004. *Veterinary Pathology* 41(1): 50-61; J. F. Soares, V. J. Chalker, K. Erles, S. Holrby, M. Waters, S. McArthur. 2004. *Journal of Zoo and Wildlife Medicine* 35(1): 25-33; A. J. Johnson, A. P. Pessier, J. F. Wellehan, R. Brown, E. R. Jacobson. 2005. *Veterinary Microbiology* 111(1-2): 107-16.