

## THE HABITAT PREFERENCES OF FISHES FROM THE LIMPOPO RIVER SYSTEM, TRANSVAAL AND MOCAMBIQUE

by

I. G. GAIGHER\*

*Abstract* - The species composition of fishes in the Limpopo River system is described. The frequency of occurrence for each habitat type is expressed as a percentage of the habitats sampled. A check list of species is presented and the species can be grouped into five habitat preferences.

### *Introduction*

The rapid industrial and agricultural development of the Transvaal is not only causing an increase in river pollution, but also necessitates the extraction of large quantities of water from the rivers, erection of weirs and dams and the transfer of water (and possibly fish species) from one river system to another. Any one, or a combination of these factors, could easily lead to the extinction of rare fish species, or of species with a specialised habitat preference.

At the moment water shortage, especially in the lower sections, presents the greatest hazard to fish species in the east-flowing rivers of the Transvaal. The most vulnerable species are those that are dependent on rapids or well-oxygenated water for existence, but all species are affected in some way or another by competition for space and food, and by being more vulnerable to diseases and predators.

It therefore became necessary to determine the status of every fish species in the Transvaal, in order to implement conservation measures if necessary. With this objective in mind, the Department of Nature Conservation, Transvaal, undertook distribution surveys in all the east-flowing rivers of the Transvaal during 1967 and 1968 (Gaigher and Pott, 1972). This paper describes the results of the determination of habitat preferences for fishes collected from the Limpopo River system during these surveys.

### *Material and Methods*

Fish were sampled at 220 different collecting sites distributed over the whole drainage area of the Limpopo River system. At each collecting site notes were made of the river type (annual or perennial), size, bottom

\* *Department of Zoology, University of Fort Hare, Alice.*

type, occurrence of aquatic plants, clearness and any other factors which could have an influence on the presence of fish at that particular site. The species of fish collected were recorded separately for each habitat type.

The determination of habitat preference was based merely on the presence or absence of species in each habitat type without considering density. Due to the selectivity of collecting gear used, it was found impractical to use abundance as an indication of habitat preference. Johnson (1967), in the determination of habitat preferences of Malaysian fresh-water fishes, came to the same conclusion, and he further found that species which have the highest frequency of occurrence for a certain habitat type, also tend to be the most abundant in that specific habitat type.

The frequency of occurrence for each habitat type, or habitat preference, is expressed as a percentage of the given habitats sampled. For example, if habitat type A is sampled twelve times, and species Z is found in six of these, then the habitat preference of species Z is 50 per cent for habitat A.

The frequency of occurrence of each fish species at five different altitudes, namely 0-305 m, 306-610 m, 611-915 m, 916-1220 m and over 1220 m was determined for each of the following habitat types:

- rapid (perennial stream);
- pool in perennial stream;
- pool in annual stream.

All the streams sampled above an altitude of 1220 m were perennial.

A more extensive classification of habitat types, including factors such as the occurrence of aquatic vegetation, bottom types, etc., was originally used. However, with the exception of a few species, these factors were found to have no marked influence on the distribution during the survey period, which was conducted during the drier months of the year.

In evaluating the results the following limitations of this method must be kept in mind:

1. The survey was done during the dry months. During the flood period fishes obviously occupy habitats that are not frequented or are not available during the dry season, in search of food, breeding sites, or in response to the natural urge to move and occupy newly available waters.
2. A few species prefer specialised habitats which do not fit into this classification. These species are discussed separately in the text.
3. Waterfalls or other barriers may isolate fish species in certain sections of the river system, thus changing its frequency of occurrence for certain altitudes where it might be able to survive. Pollution will have the same influence on the results.

### *Results and Discussion*

A check list of fishes collected from the Limpopo River System

#### FAMILY LEPIDOSIRENIDAE

*Protopterus annectens brieni* Poll, 1961.

#### FAMILY ANGUILLIDAE

*Anguilla marmorata* Quoy & Gaimard, 1824.

*A. nebulosa labiata* Peters, 1852.

*A. mossambica* Peters, 1852.

*A. bicolor bicolor* McClelland, 1845.

#### FAMILY MORMYRIDAE

*Petrocephalus catostoma* (Günther), 1866.

*Gnathonemus macrolepidotus* (Peters), 1852.

#### FAMILY CHARACIDAE

*Hydrocynus vittatus* Castelnau, 1861.

*Alestes imberi* Peters, 1852.

*Micralestes acutidens* (Peters), 1852.

#### FAMILY CYPRINIDAE

*Barbus polylepis* Boulenger, 1907.

*B. mattozi* Guimaraes, 1884.

*B. paludinosus* Peters, 1852.

*B. eutaenia* Boulenger, 1904.

*B. unitaeniatus* Günther, 1866.

*B. neefi* Greenwood, 1962.

*B. viviparus*, M. Weber, 1897.

*B. toppini* Boulenger, 1916.

*Labeo rubropunctatus* Gilchrist & Thompson, 1913.

*L. rosae* Steindachner, 1894.

*L. molybdinus* du Plessis, 1963.

*L. cylindricus* Peters, 1852.

*L. ruddi* Boulenger, 1907.

*B. marequensis* A. Smith, 1841.

*B. trimaculatus* Peters, 1852.

*B. afrohamiltoni* Crass, 1960.

*B. anoplus* M. Weber, 1897.

*B. lineomaculatus* Boulenger, 1903.

*B. treuensis* Groenewald, 1958.

*B. annectens* Gilchrist & Thompson, 1917.

*Beirabarus radiatus* (Peters), 1853.

*Barilius zambezensis* (Peters), 1852.

*Engraulicypris brevianalis* (Boulenger), 1908.

#### FAMILY SCHILBEIDAE

*Eutropius depressirostris* (Peters), 1852.

#### FAMILY AMPHILIIDAE

*Amphilius platyichir* Günther, 1864.

#### FAMILY CLARIIDAE

*Clarias gariepinus* (Burchell), 1822. *C. ngamensis* Castelnau, 1861.

#### FAMILY MOCHOKIDAE

*Synodontis zambezensis* Peters, 1852. *C. swierstrai* v. d. Horst, 1931.

*Chiloglanis paratus* Crass, 1960. *C. pretoriae* v. d. Horst, 1931.

## FAMILY CYPRINODONTIDAE

*Aplocheilichthys johnstonii* (Günther), 1893. *A. katangae* (Boulenger), 1912.

## FAMILY CICHLIDAE

*Tilapia mossambica* Peters, 1852. *T. sparrmanii* A. Smith, 1840.  
*T. melanopleura* Dumeril, 1859. *Chetia flaviventris* Trewavas, 1961.  
*Haplochromis brevis* (Jubb), 1968.  
*Serranochromis meridianus* Jubb, 1967.  
*Hemihaplochromis philander* M. Weber, 1897.

## FAMILY GOBIIDAE

*Platygobius aeneofuscus* (Peters), 1852.  
*Glossogobius giuris* (Hamilton-Buchanan), 1822.

### *Habitat Preference*

Based on habitat preference and distribution, the fishes of the Limpopo River system can be divided into five ecological groups, namely:

- A. Species with a wide distribution (high- and lowveld) and a wide habitat preference;
- B. Species with a wide habitat preference, but which are confined to the warmer middle- and lowveld streams;
- C. Pool-loving species confined to the warmer middle- and lowveld streams;
- D. Species confined to well-oxygenated middle- and highveld streams;
- E. Species with a restricted distribution and/or habitat preference.

### Group A: (Fig. 1)

Unspecialised species with a wide distribution. *B. marequensis* is included in this group because it is so widespread and abundant. It differs from the other species, in being dependent on rapids for breeding purposes, and is therefore found mainly in perennial streams. It is absent from the lower section of the Limpopo River in Mocambique which is devoid of rapids. Although widespread, *T. sparrmanii* is specialised to a certain extent in that it prefers habitats with abundant rooted aquatic vegetation.

Group A includes the following species:

*Barbus marequensis*  
*B. trimaculatus*  
*B. paludinosus*  
*Clarias gariepinus*  
*Tilapia sparrmanii*  
*Hemihaplochromis philander*




Fig. 1. The percentage frequencies of occurrence at different altitudes of group A fishes in the Limpopo River system, 1968.




Fig. 2. The percentage frequencies of occurrence at different altitudes of group B fishes in the Limpopo River system, 1968.

Group B: (Fig. 2)

These species are unspecialised and also abundant, but differ from the previous group in being absent from the coldest highveld streams (this difference is not so apparent from the figures, but is clearly shown in the distribution maps).

*Labeo cylindricus* and *L. molybdinus* were collected mainly from rapids, but both are abundant in pans and river sections in the lowlands of Mocambique, and are therefore not dependent on rapids for existence. The results show that they prefer rapids, most probably for feeding purposes, if these are available.

Group B includes the following species:

- Micralestes acutidens*
- Barbus unitaeniatus*
- B. viviparus*
- Engraulicypris brevianalis*
- Labeo cylindricus*
- L. molybdinus*

Group C: (Fig. 3)

Pool-loving species confined to the warmer middle- and lowveld streams. They are not dependent on rapids for feeding or breeding purposes and all do well in irrigation dams. Although not strictly freshwater fishes, *Glossogobius giuris* and *Platygobius aeneofuscus* can be added to this list as far as distribution and habitat preference in the Limpopo River system is concerned.

Group C includes the following species:

- Petrocephalus catostoma*
- Gnathonemus macrolepidotus*
- Alestes imberi*
- Barbus mattozi*
- B. afrohamiltoni*
- B. annectens*
- B. toppini*
- Beirabarus radiatus*
- Labeo rubropunctatus*
- L. rosae*
- L. ruddi*
- Eutropius depressirostris*
- Synodontis zambezensis*
- Tilapia mossambica*
- T. melanopleura*











Fig. 3. The percentage frequencies of occurrence at different altitudes of group C fishes in the Limpopo River system, 1968.





■ RAPID (PERENNIAL STREAM)  
 ▨ POOL IN PERENNIAL STREAM  
 □ POOL IN ANNUAL STREAM

Group D: (Fig. 4)

These species are confined to perennial middle- and highveld streams. They are less abundant than the previous groups, but are not so specialised that they are in any danger of disappearing from the system. Only *Chetia flaviventris* is endemic to the Limpopo River system. Although this species does not have a wide distribution in the system, it is well adapted to conditions in irrigation dams. In certain dams, for example, the Hartebeestpoort Dam near Pretoria and Buffelspoort Dam near Rustenburg, it has become so abundant that it is a nuisance to anglers, and the numbers must be controlled from time to time.




Fig. 4. The percentage frequencies of occurrence at different altitudes of group D fishes in the Limpopo River system, 1968.

Group D includes the following species:

- Barbus polylepis*
- B. eutaenia*
- B. anoplus*
- B. neefi*
- B. lineomaculatus*
- Chetia flaviventris*

Group E: (Figs. 5 and 6)

The following fish species have a restricted distribution and/or specialised habitat preference and will therefore be easily affected by water shortage or other habitat changes.

1. *Protopterus annectens brieni*

This interesting species is confined to floodplains and pans that dry up wholly or partially during the dry months. It is abundant along the lower part of the Limpopo River in Mocambique. The future erection of irrigation dams in the upper reaches that are large enough to prevent inundation of flood plains and pans during the summer months, could possibly lead to the disappearance of the lungfish in this area.

2. *Hydrocymus vittatus*

Tiger fish are abundant in Mocambique, but in the Transvaal they are virtually confined to the Kruger National Park, where they are not available to anglers. Water shortage and the erection of weirs are probably the main reasons for the disappearance of this fine angling fish from higher stretches of the system where it previously occurred.

3. *Barbus treurenensis*

This minnow is endemic to the upper reaches of the Blyde River, Eastern Transvaal. For a long time it was only known from one tributary, the Treur River. The introduction of trout and bass to the Treur River has probably caused its disappearance and no specimens could be found during this survey despite an intensive search. Subsequently another population has been found by the Provincial Fisheries Institute, Lydenburg in a tributary upstream of the Treur River where attempts are being made to provide a sanctuary for this rare species (Jubb, 1972).

4. *Barilius zambezensis*

This colourful little fish is confined to clear, fast-flowing perennial middle- and lowveld streams. It was never found in standing water. Several middle- and lowveld stretches of previously perennial streams have become annual during the past few years. If this trend increases, it could possibly lead to the disappearance of *B. zambezensis* from the Limpopo River system.

5. *Amphilius platyichir*, *Chiloglanis paratus* and *C. pretoriae*

Although these species have a wide distribution in the Limpopo River system, they have a highly specialised habitat preference in being virtually confined to rapids. They might therefore disappear from sections of the river system where flow ceases, even only for a short period.

## 6. *Chiloganis swierstrai*

This species is confined to fast-flowing sections of middle- and lowveld streams with a sandy or pebbly bottom. This type of habitat is rare in the Limpopo River system, especially during the dry months when the water tends to flow through the sand substratum instead of over it.

During this survey *C. swierstrai* was only found in the Steelpoort and Olifants Rivers. During surveys done before 1965 by the Provincial




Fig. 5. The percentage frequencies of occurrence at different altitudes of group E fishes in the Limpopo River system, 1968.




Fig. 6. The recorded distributions of *Barbus treurenis*, *Chiloglanis swierstrai*, *Aplocheilichthys spp.*, *Haplochromis brevis* and *Serranochromis meridianus* in the Limpopo River system, 1968.

Fisheries Institute, it was also found in the Letaba and Limpopo Rivers, and Pienaar (1968) collected it from four different sites in the Letaba River.

7. *Clarias ngamensis* has a limited distribution in the lower parts of the system in Mocambique, where it is abundant. The reasons for its absence from the higher reaches are unknown.

7. *Clarias ngamensis* and *C. gariepinus* have a limited distribution in the lower parts of the system in Mocambique, where they are abundant. The reasons for its absence from the higher reaches are unknown.

8. *Aplocheilichthys* spp.

These two species are eurotherm, but are confined to clear pools and slow-flowing sections of perennial streams with abundant aquatic vegetation. *A. katangae* was only found in the Matlapatzi River, a small tributary of the Olifants River (Fig. 6). This is its only known locality in the Transvaal. *A. johnstonii* is abundant in the lower parts of the system, especially in the floodplains and pans in Mocambique. In the Transvaal it has a scattered distribution in the upper reaches of the Mogalakwena, Mogol and Marico Rivers.

9. *Haplochromis brevis* and *Serranochromis meridianus*

These cichlids are endemic to the Incomati and Limpopo River systems. During this survey they were only found in coastal pans adjacent to the Limpopo River (Fig. 6).

#### *Acknowledgements*

The author is greatly indebted to the following: The Director of Nature Conservation, Transvaal, who enabled him to do the survey while at the Provincial Fisheries Institute, Lydenburg, Republic of South Africa; Dr. R. A. Jubb of the Albany Museum for fish identifications; Dr. U. de V. Pienaar of the National Parks Board for assistance and advice during surveys in the Kruger National Park and Mr. F. J. van der Merwe of the Provincial Fisheries Institute who assisted with the surveys.

#### REFERENCES

- GAIGHER, I. G. and R. McC. POTT. 1972. A check list of indigenous fish in the east-flowing rivers of the Transvaal. *Limn. Soc. S.A. News Letter* 18: 26-32.
- JOHNSON, D. S. 1967. Distributional patterns in Malayan freshwater fish. *Ecology* 48(5): 722-730.
- JUBB, R. A. 1972. Disappearing river fishes. *Afr. wild L.*, 26(2): 48-52.
- PIENAAR, U. DE V. 1968. The freshwater fishes of the Kruger National Park. *Koedoe* 11: 1-82.