

Adjustable Strabismus Surgery: An Early Glance

Sana Nadeem, B.A Naeem, Farman Khan

Pak J Ophthalmol 2018, Vol. 34, No. 2

See end of article for authors affiliations

Correspondence to:
Sana Nadeem
Eye Department
Fauji Foundation Hospital,
Rawalpindi
Email:
sana.nadeem018@gmail.com

Purpose: To assess the short term success rate of adjustable suture technique on rectus muscle strabismus surgery in terms of postoperative alignment

Study Design: Prospective, interventional study

Place and Duration of Study: Eye Department, Fauji Foundation Hospital, Rawalpindi, from 25th June, 2016 to 25th December, 2017

Material and Methods: We carried out a study using the fornix approach for adjustable squint surgery, in mainly horizontal strabismus in adults and cooperative children, to finely tune the postoperative alignment. The preoperative deviation, strabismus type, patterns, were measured and analyzed. The early postoperative alignment was measured at 6 weeks postoperatively, to assess the success of the adjustable suture technique. A sliding noose knot was used to adjust the sutures, 1 hour postoperatively under topical proparacaine anesthesia, after the effects of general anesthesia had worn off.

Results: 31 patients were included in this study. The majority were female being 23 (74.5%). The age ranged from 9 to 37 years, with the mean age 16.87 ± 5.5 years. All consenting adults and teenagers, with strabismus were included in this study and operated via the adjustable suture technique. The mean preoperative deviation was $49.38 \pm 16.29^{\Delta}$ prism diopters, and the mean postoperative deviation was $3.5 \pm 5.42^{\Delta}$ prism diopters. The difference between the two was statistically significant ($p < 0.05$), using the Wilcoxon Signed Ranks test ($p = 0.000$). Early surgical success defined as alignment within $\pm 10^{\Delta}$ (prism diopters) of orthotropia at the end of 6 weeks after surgery, was found in 27 (87.1%) of our strabismus cases.

Conclusion: Adjustable strabismus surgery is associated with excellent short-term postoperative outcomes in terms of alignment and patient satisfaction. Adjustable sutures should be considered in all strabismus cases, whether adults or children.

Key Words: short term, success, adjustable, strabismus, fornix approach, alignment.

Adjustable strabismus surgery¹ is an art, dating back to 1907, with the ideation of Bielchowsky, O'Connor and Harms who first introduced this technique to refine postoperative alignment. Jampolsky¹ later re-introduced the adjustable suture technique to achieve stable motor alignment and sensory improvement. Although most strabismus surgeons who use adjustable sutures,

including me, would prefer them for all their patients regardless of the cause of strabismus, or age of the patient; there are certain indications² where they are necessary for good cosmetic results: restrictive strabismus including thyroid myopathy, anesthetic myotoxicity, scleral buckles, paralytic strabismus, and diplopia to name a few. This art however, has failed to gain universal acceptance so far, and especially in

Pakistan³. Strabismus surgery can be tricky³ in terms of its outcomes, despite accurate measurements and nomograms followed, and may yield different results in different patients, in spite of the same amount of alignment. Thus, the need arises, to adjust the eyes postoperatively, to maximize the chances of success. Adjustable sutures give the operating surgeon a 'second chance' at achieving a stable alignment⁴.

In June 2016, we started doing adjustable strabismus surgery, and found it to achieve superior cosmesis, and improve binocular vision and diplopia⁵. Since we are relatively new at this technique, we decided to review and share our early postoperative alignment at six weeks⁶ postoperatively, as this is the time, which reflects the chance of the patient achieving eventual long term stable alignment of eyes.

The *aim* of this study is to describe the adjustable suture technique and to assess its effectiveness and success in the early postoperative period, prospectively, in addition to highlight the importance and benefits of postoperative adjustment on postoperative outcomes. The importance of sharing our experiences with others is to benefit those who want to help their patients in achieving the best postoperative cosmetic results.

MATERIAL AND METHODS

All consenting adults and co-operative teenagers with strabismus were included in this ongoing study, reaching 31 patients. This study was carried out in the Department of Ophthalmology, Fauji Foundation Hospital, Rawalpindi, which is a tertiary care, teaching hospital affiliated with Foundation University Medical College; from 25th June 2016 to 25th December 2017. Approval from the ethical committee was taken. The strabismus cases included both horizontal and vertical strabismus, with only one horizontal muscle being used for adjustment. Patients with previous history of strabismus surgery were also included. The first author performed all surgeries. Restrictive strabismus, myasthenia gravis, and uncooperative children less than 9 years of age were excluded.

Visual acuity was documented for every case with a refractive correction given to patients prior to consideration for surgery. The type of strabismus was noted for each patient. The preoperative angle of deviation was assessed by the prism cover test for both near and distance with the refractive correction in place. In certain cases of sensory strabismus with poor fixation, the Krimsky test was used for analysis of the

angle or a pen torch used as a target for near and distance. The angle of deviation was measured for both near and distance as well as in up gaze, downgaze, right and left gaze. However, the distance angle with refractive correction in place was considered as the angle of deviation in all cases, and the surgical alignment was sought to correct this angle. At the time of suture adjustment, though, both near and distance alignment was corrected. Exception to this was accommodative refractive esotropia, for which the near deviation with distance spectacles in place was considered for correction of the alignment. A plus sign (+) was assigned to an exotropic angle and a negative sign (-) was assigned to an esotropic angle. The measurements were taken by the operating surgeon and a certified orthoptist, and repeated by the operating surgeon one day prior to surgery, to obtain maximum cosmesis. Binocular vision and stereopsis were assessed by the Titmus fly test and Worth four-dot test, routinely by the orthoptist preoperatively. A thorough eye examination was performed including fundus and intraocular pressures, and was documented.

All surgeries were performed under general anesthesia. A drop of phenylephrine 10% (Ethifrin®) were instilled into the conjunctival fornix at the beginning of surgery in each eye. The fornix approach for strabismus surgery was used in every case. Each muscle was hooked, and then secured with a double armed 6-0 vicryl (polyglactin 910) absorbable suture, which was passed through the sclera at its insertion, or transposed above or below the insertion in case of 'A' or 'V' patterns, in a 'hang-back'^{7,8} fashion. The medial recti were transposed towards the apex, and the lateral recti were transposed away from the apex, in case of 'A' or 'V' patterns. The recessed muscles were mostly placed for adjustable purpose, with the required recession held in place by Guyton's⁹ modification of the sliding noose knot, which was fashioned with a 6-0 Vicryl suture. The amount of 'hang-back' recession was calculated for each patient using standard tables^{4,10,11}. The traction suture for holding the sclera for postoperative adjustment was created with ethibond 5-0 in every case. For the non-adjustable recessions, the muscle was tied and allowed to 'hang-back' from its insertion, with the required amount of recession calculated as required. Resections were also put up for adjustment in one case only, but avoided mostly, as they tend to cause more pain. For resections, the amount of resection is overcorrected by 2 mm, and allowed to 'hang-back' for this distance, to

be adjusted if required postoperatively. Only one muscle was kept on an adjustable sliding noose knot per case.

All patients were assessed for alignment and final adjustment at least 1 hour or more after surgery, in the recovery room, to allow time for the effects of general anesthesia to wear off¹². The eyes were anesthetized topically with Alcaine® (proparacaine hydrochloride 0.5%) eye drops. The patients were assessed with the cover-uncover test at distance and near, with a torch light for distance if the vision was blurred (due to viscoelastic, antibiotic/steroid drops placed postoperatively or pupillary dilatation with phenylephrine or general anesthesia), or a distance readable target, and for near an accommodative target was used. If the alignment was satisfactory, with no movement on cover testing, the sutures were tied off in their existing position, held in place by the sliding noose, which was removable after tying the ends of the muscle sutures. Thereafter, the traction knot was cut, and the conjunctiva was sutured with at least one 6-0 vicryl suture. The final tying off point was orthotropia or maximum possible under-correction as required. In cases of exotropia, the goal was orthotropia or mild esotropia. In cases of esotropia, the goal was either orthotropia, if achieved, or slight under-correction.

The alignment was noted postoperatively the next day, at 2 weeks and 6 weeks after surgery. The patients were given postoperative topical steroid and antibiotic drops twice a day and ointment at night for a minimum of 2 weeks. The follow up is being continued to assess long term postoperative alignment as well.

Early surgical success was defined as alignment within $\pm 10^A$ (prism diopters) of orthotropia at the end of 6 weeks after surgery. This postoperative residual deviation was the average of the distance and near deviations noted on prism cover testing. Although both horizontal, vertical and complex strabismus were included in our study, the horizontal alignments preoperatively and postoperatively, were mainly assessed and analyzed for surgical success purpose.

The results were noted, tabulated and analyzed using the SPSS statistics version 20. Frequencies were calculated for age, gender, type and pattern of strabismus, surgical procedure performed, as well as the follow-up. Statistical analysis of success rate was done, and assessment of statistically significant differences between the preoperative and

postoperative strabismus deviations was analyzed by the *Wilcoxon Signed Ranks test*. The success rate was compared and analyzed for the type of strabismus as well.

RESULTS

31 patients were included in this study, with predominantly 23 (74.2%) females and 8 (25.8%) males. The mean age was 16.87 ± 5.5 years with a range from 9 years to 37 years. The early postoperative alignment was measured at 6 weeks follow up, and the average follow up was 52.7 ± 13.76 days, with an actual range from 38 to 94 days.

The patients were classified based on type of deviation and the majority of them were exotropic with 16 (51.6 %) cases [Table 1]. Table 1 categorizes the patients based on type of strabismus, with predominance of purely horizontal cases in 17 (54.8 %) cases; the rest being a combination of horizontal, vertical and/or complex strabismus. Associated patterns, A, V, X, Y were tabulated in Table 2, with predominance of V-pattern.

Table 1: Type of Strabismus.

Deviation Type	Frequency (Percent)
Exotropia	16 (51.6)
Esotropia	6 (19.4)
Exotropia & DVD ^φ	2 (6.5)
Esotropia & DVD ^φ	1 (3.2)
Exotropia & Hypertropia	5 (16.1)
Esotropia & Hypertropia	1 (3.2)
Horizontal	17 (54.8)
Horizontal and Vertical	7 (22.6)
Horizontal & Complex [€]	4 (12.9)
Horizontal, Vertical & Complex [€]	3 (9.7)

^φ Dissociated vertical deviation

[€] Sensory, paralytic strabismus or DVD

Table 2: Pattern of Strabismus if Present.

Pattern	Frequency (Percent)
V-pattern	20 (64.5)
A-pattern	4 (12.9)
None	7 (22.6)

Table 3: Surgical Procedure Performed.

Surgical Procedure	Frequency (Percent)
BLRc¶	12 (38.7)
BMRc§	5 (16.1)
BMRs¥	1 (3.2)
MRc ^α + LRs [×]	1(3.2)
MRs ^ø + LRC ^H	6 (19.4)
BLRc¶ + MRs ^ø	4 (12.9)
BMRc§ + LRs [×]	1 (3.2)
MRc ^α	1 (3.2)

- ¶ Bilateral recessions
- § Bimedial recessions
- ¥ Bimedial resections
- Unilateral medial rectus recession
- × Unilateral lateral rectus resection
- ø Unilateral medial rectus resection
- H Unilateral lateral rectus recession

The various surgical procedures performed to correct horizontal component of strabismus are listed in Table 3. Associated vertical deviations were addressed with concurrent procedures on the vertical Recti or obliques, which are listed in Table 4. A summary of the data in accordance with the type of deviation is elaborated in Table 5.

The mean preoperative deviation was 49.38 ± 16.29^Δ (prism diopters), with a range from 23 to 85^Δ. The mean postoperative deviation was 3.5 ± 5.42^Δ

(prism diopters), with a range from zero to 20^Δ. The difference between the preoperative and postoperative deviation was analyzed by the *Wilcoxon Signed Ranks test* and found to be statistically significant (p = 0.000) [Table 6]. Early surgical success defined as satisfactory alignment within ± 10^Δ (prism diopters) of orthotropia at the end of 6 weeks after surgery, was found in 27 (87.1%) of our strabismus cases, and were tabulated for each deviation type [Table 7], and under corrections were found in 3 (9.6%) cases of exotropia and 1 (3.2%) case of esotropia only. Figures 1-3 depict our post-operative success. Reoperation has not been required in any of these cases so far. No complication was encountered in any patient during suture adjustment and all proceeded smoothly with minimal patient discomfort.

Table 4: Surgery on Vertical Recti & Obliques.

Surgery on Vertical Recti & Obliques	Frequency (Percent)
Unilateral IO ^α myectomy	3 (9.7)
Bilateral IO ^α myectomies	6 (19.4)
IO ^α anteriorization	1 (3.2)
SR ^β transposition	1 (3.2)
Bilateral IO ^α myectomies + SRC ^Σ	1 (3.2)
None	19 (61.3)

- α Inferior oblique
- β Superior rectus
- Σ Superior rectus recession

Table 5: Data Analysis According to Deviation Type.

Type of Deviation	Age in Years (mean ± SD)	Gender		Preoperative Angle (PD) (Mean ± SD) ^Ω	Postoperative Angle (PD) (Mean ± SD)	Pattern	
		Male	Female			V-Pattern	A-Pattern
Exotropia	19.06 ± 6.18	4	12	48.1 ± 16.7	4.46 ± 6.77	9	2
Esotropia	14.17 ± 4.4	2	4	60.1 ± 13.3	2.58 ± 4.34	4	2
Exotropia & DVD ^ϕ	13.5 ± 4.9	1	1	34 ± 8.4	1.00 ± 1.41	2	0
Esotropia & DVD ^ϕ	15.0	1	0	70	0	1	0
Exotropia & Hypertropia	15.0 ± 3.53	0	5	47.8 ± 10.7	2.30 ± 2.58	4	0
Esotropia & Hypertropia	16.0	0	1	23	8 ± 3.50	0	0

- PD Prism diopters
- SD Standard deviation

Ω distance deviation in all cases except accommodative esotropia in which near deviation with distance correction was taken

Table 6: Preoperative and Postoperative Deviations.

Descriptive Statistics	Mean	Std. Deviation	Minimum	Maximum	Percentiles		
					25th	50th (Median)	75th
Preoperative Deviation (PD)	49.3871	16.29249	23.00	85.00	40.0000	45.0000	65.0000
Postoperative Deviation (PD) at 6 weeks	3.5000	5.42525	.00	20.00	.0000	1.0000	6.0000

PD Prism diopters

Test Statistics^a

	Postoperative Deviation at 6 weeks - Preoperative Deviation (prism diopters)
Z	-4.862 ^b
Asymp. Sig. (2-tailed)	.000

- a. Wilcoxon Signed Ranks Test
- b. Based on positive ranks.

Table 7: Surgical Success.

		Surgical Success with Postoperative Deviation within 10 Prism Diopters	
		Yes	No
Total	31	27 (87.1)	4 (12.9)
Deviation Type	Exotropia	13 (41.9)	3(9.6)
	Esotropia	5 (16.1)	1 (3.2)
	Exotropia & DVD ^φ	2 (6.4)	0
	Esotropia & DVD ^φ	1 (3.2)	0
	Exotropia & Hypertropia	5 (16.1)	0
	Esotropia & Hypertropia	1 (3.2)	0

^φ Dissociated vertical deviation



Fig. 1 (A): 20 year old girl with congenital alternate esotropia of 70Δ, a left hypertropia of 8Δ, and a V-pattern of 15Δ. Bimedial recessions of 7mm OD and 8 mm OS after adjustment and a left inferior oblique myectomy was done.



Fig. 1 (B): Postoperatively, she is well aligned.




Fig. 2 (A): A 15-year-old girl with an alternate exotropia of 70Δ and a V-pattern of 32Δ was managed by bilateral lateral rectus recessions of 7mm and a right medial rectus resection of 6 mm, and bilateral inferior oblique myectomies. No adjustment was needed postoperatively and the sutures were tied off.




Fig. 2 (B): Postoperatively she is orthotropic.



Fig. 3 (A): A 12 year old boy with an alternate exotropia of 40Δ and a V-pattern of 20Δ was managed with bilateral lateral rectus recessions 6mm OD and 8 mm OS after adjustment, along with a half tendon width upward transposition of both muscles was done.



Fig. 3 (B): Postoperatively, he is orthotropic, with resolution of the V-pattern.

DISCUSSION

In our study, we found adjustable strabismus surgery to be highly effective for postoperative satisfactory alignment with a high success rate of 87.1%. The patients experienced no adverse effects or complications during the adjustable procedure, and did not report much discomfort during suture tying. We minimized patient discomfort by putting only one muscle on an adjustable noose knot per procedure, and this too has borne fruitful results. Suture adjustment was needed in the majority of patients postoperatively, amounting to 26 (83.8%) cases, and no adjustment was required for 5 (16.1%) cases, thus indicating the need to fine tune the postoperative alignment to achieve orthotropia and better surgical outcomes, because the nomograms and tables may not be appropriate for each individual case.

Thus, our study findings indicating positive and encouraging outcomes with the adjustable suture technique are consistent with numerous studies worldwide, which report high success rates with the likes of Wisnicki, Repka and Guyton¹³, who reported a huge success rate of this adjustable procedure in a massive 290 patients, and rate of reoperation in just 9.7%. Eino et al¹⁴, reported a success rate of 91.7% in 109 patients, and Tripathi et al¹⁵, observed a higher success rate with adjustable versus non-adjustable surgery in 443 total cases. Engel et al¹⁶ reported a high short-term success rate of 88% in their adjustable suture technique in 61 children, and so did Awadein and Guyton¹², with their study on children and infants using propofol anesthesia, who noted a success rate of 79%. Nihalani and Hunter¹⁶ also noted a high success rate for both vertical and horizontal muscle surgery

using a short tag noose technique. Budning et al¹⁷ developed a short adjustable suture with high success in 304 patients. Locally, a study done by Shakir et al¹⁸ at LRBT Free Base Eye Hospital in Karachi, reported a success rate of 88% in 18 exotropic subjects with adjustable technique.

Several other authors Park et al¹⁹ and Bishop et al²⁰, have contrasting evidence in their reported studies, where they did not find any difference in surgical success, when comparing adjustable versus non-adjustable techniques.

The strengths of our study are its success in terms of surgical results. There are both vertical and horizontal strabismus patients included along with complex cases like consecutive strabismus, DVD, patterns, and one case of a sixth nerve palsy. Transpositions were also performed during surgery on these patients. The surgeries were carried out in a single step to restore ocular alignment, rather than splitting the large alignments at a second stage.

A few limitations of our study are a relatively small sample size, because we wanted to share our early postoperative results, and we do intend to add more patients to this study with time. Strabismus patients are not that frequent in our set up and using this technique on more patients is our priority, for the interest of the patients. Four of our patients did not have a successful postoperative alignment, and we could attribute it to measurement errors^{21,22}, or muscle abnormalities, hypoplasia or pulley anomalies, which can be assessed preoperatively with neuroimaging²³. Also early alignment may vary from subsequent long-term alignment, which is more important to the patient especially, and to the operating surgeon, although we have observed them to be quite similar in many of the cases. In addition, we used only one muscle per patient for the adjustable suture technique both to avoid excessive discomfort, and to save time. Others² recommend using adjustable sutures on all muscles, except on the inferior oblique, in order to adjust the muscles symmetrically and achieve a balanced alignment between the two eyes. We intend to broaden our experience on adjustable strabismus, to include all cases including children, to offer our patients a better chance of success. Some patients have not returned for a follow up, which we had to exclude from the study, decreasing our eventual total number of cases.

Future work required is a large-scale study with more subjects, which we are recruiting, and

assessment of long-term outcomes at one year postoperatively and beyond. Using the adjustable suture on multiple or all muscles may further enhance the success rate of this procedure. Infants and children¹² also need to be given a chance at better postoperative success with the use of propofol anesthesia. Superior oblique adjustment can also be done with this technique, if the need arises to operate on this muscle.

The need arises to encourage strabismus surgeons, to use this adjustable technique to maximize the chances of their surgical success, and to reduce the number of reoperations needed to satisfy the patients.

CONCLUSION

Adjustable strabismus surgery has been found to yield excellent short-term postoperative outcomes in terms of alignment and patient satisfaction. The adjustable suture technique should be considered in all strabismus cases, whether horizontal, vertical or cyclovertical; simple or complex strabismus, and irrespective of the subjects being adults or children.

Author's Affiliation

Dr. Sana Nadeem
Assistant professor
Ophthalmology Department
Fauji Foundation Hospital/FUMC, Rawalpindi

Prof. Dr. B.A Naeem
Professor and Head
Ophthalmology Department
Fauji Foundation Hospital/FUMC, Rawalpindi

Farman Khan
Orthoptist
Ophthalmology Department
Fauji Foundation Hospital/FUMC, Rawalpindi

Role of Authors

Dr. Sana Nadeem
Principle author, Performed all surgeries, Orthoptic assessment & Evaluation of all patients, Data Analysis.

Prof. Dr. B.A Naeem
Reviewing author, Opinion about patient management, Head of Department

Farman Khan
Orthoptic assessment & Fusion of all the patients

REFERENCES

1. **Vasconcelos GC, Guyton DL.** Adjustable sutures in strabismus surgery: why surgeons either love them or hate them after three decades. Editorial. *Arc Bras Oftalmol.* 2014 Aug; 77 (4): 5-6.
2. **Guyton DL.** Adjustable sutures. In: *Clinical Atlas of Procedures in Ophthalmic and Oculofacial surgery.* Second Ed. Oxford university press: USA, 2011: P 592.
3. **Nihalani BR, Hunter DG.** Adjustable suture strabismus surgery. *Eye.* 2011 Oct; 25 (10): 1262-1276.
4. **Coats DC, Olitsky SE.** Strabismus surgery and its complications. Springer: Berlin, 2007: P 37-39.
5. **Ferdi A, Kelly R, Logan P, Dooley I.** Outcomes of adjustable strabismus surgery an Irish University Hospital. *Int Ophthalmol.* 2016 Nov 8; 37 (5): 1215-1219.
6. **Liebermann L, Hatt SR, Leske DA, Holmes JM.** Adjustment versus no adjustment when using adjustable sutures in strabismus surgery. *J AAPOS.* 2013 Feb; 17 (1): 38-42.
7. **Capó H, Repka MX, Guyton DL.** Hang-back lateral rectus recessions for exotropia. *J Pediatr Ophthalmol Strabismus,* 1989; 26: 31-4.
8. **Repka MX, Guyton DL.** Comparison of hang-backmedial rectus recession with conventional recession. *Ophthalmology,* 1988; 95: 782-7.
9. **Deschler EK, Irsch K, Guyton KL, Guyton DL.** A new, removable, sliding noose for adjustable-suture strabismus surgery. *J AAPOS.* 2013 Oct; 17 (5): 524-7.
10. **Lueder GT, Archer SM, Hered RW, Karr DJ, Kodsi SR, Kraft SP, et al.** Pediatric Ophthalmology and Strabismus. Section 6. Basic and Clinical Science Course. American Academy of Ophthalmology. San Francisco, 2015-2016: p 182.
11. **Yanoff M, Duker JS.** *Ophthalmology.* Third Edition. Mosby: St. Louis, 2009: p 1331-1338.
12. **Awadein A, Sharma M, Bazemore GM, Saeed HA, Guyton DL.** Adjustable suture strabismus surgery in infants and children. *J AAPOS.* 2008; 12: 585-590.
13. **Wisnicki HJ, Repka MX, Guyton DL.** Reoperation rate in adjustable strabismus surgery. *J Pediatr Ophthalmol Strabismus,* 1988; 25: 112-114.
14. **Eino D, Kraft SP.** Postoperative drifts after adjustable suture strabismus surgery. *Can J Ophthalmol.* 1997; 32: 163-169.
15. **Tripathi A, Haslett R, Marsh IB.** Strabismus surgery: Adjustable good for all? *Eye,* 2003; 17: 739-742.
16. **Nihalani BR, Whitman MC, Salgado CM, Loudon SE, Hunter DG.** Short tag noose technique for optional and late suture adjustment in strabismus surgery. *Arch Ophthalmol.* 2009 Dec; 127 (12): 1584-1590.
17. **Budning AS, Day C, Nguyen A.** The short adjustable suture. *Can J Ophthalmol.* 2010; 45: 359-362.
18. **Shakir M, Kamil Z, Zafar S, Bokhari SA, Rizvi F.** Adjustable sutures in constant exotropia. *Pak J Ophthalmol.* 2013; 29 (4): 192-196.
19. **Park YC, Chun BY, Kwon JY.** Comparison of the stability of postoperative alignment in sensory exotropia: adjustable versus non-adjustable surgery. *Korean J Ophthalmol.* 2009; 23: 277-280.
20. **Bishop F, Doran RM.** Adjustable and non-adjustable strabismus surgery: a retrospective case-matched study. *Strabismus,* 2004; 12: 3-11.
21. **Pediatric Eye Disease Investigator Group.** Inter-observer reliability of the prism and alternate cover test in children with esotropia. *Arch Ophthalmol.* 2009 Jan; 127 (1): 59-65.
22. **Thompson JT, Guyton DL.** Ophthalmic Prisms. Measurement errors and how to minimize them. *Ophthalmology,* 1983 Mar; 90 (3): 204-210.
23. **Patel JR, Gunton KB.** The role of imaging in strabismus. *Curr Opin Ophthalmol.* 2017 Sep; 28 (5): 465-469.