

1 SUBMITTED 8 DEC 21  
2 REVISIONS REQ. 21 FEB 22; REVISIONS RECD. 5 APR 22  
3 ACCEPTED 19 APR 22  
4 **ONLINE-FIRST: APRIL 2022**  
5 DOI: <https://doi.org/10.18295/squmj.4.2022.034>

6

7 **Recurrent scrotal Arteriovenous Malformation as a Slowly**  
8 **Increasing Left Testicular Swelling**

9 *A case report*

10 **Faisal M. Alashaikh,<sup>1,2</sup> \*Moustafa S. Alhamadh,<sup>1,2</sup> Sulaiman I.**  
11 **Alsugair,<sup>1,2</sup> Abdullallah Alkhayal,<sup>1,2,3</sup> Saad Abumelha,<sup>1,2,3</sup> Yousof Al**  
12 **Zahrani,<sup>2,4</sup> AbdulRahman BinSwilim<sup>5</sup>**

13

14 <sup>1</sup>College of Medicine, King Saud ben Abdulaziz University for Health Sciences  
15 (KSAU-HS), Ministry of the National Guard-Health Affairs, Riyadh, Saudi Arabia;

16 <sup>2</sup>King Abdullah International Medical Research Center, Ministry of the National  
17 Guard-Health Affairs, Riyadh, Kingdom of Saudi Arabia; <sup>3</sup>Department of Surgery,  
18 Division of Urology and <sup>4</sup>Department of Medical Imaging, Division of Interventional  
19 Radiology, King Abdulaziz Medical City, Ministry of the National Guard-Health  
20 Affairs, Riyadh, Kingdom of Saudi Arabia; <sup>5</sup>Department of Medical Imaging, King  
21 Khalid University Hospital, Riyadh, Kingdom of Saudi Arabia.

22 \*Corresponding Author's e-mail: Alhamadhmo@gmail.com

23

24 **Abstract**

25 Arteriovenous malformations (AVMs) are benign vascular lesions. Although, the  
26 majority of AVMs occur in the central nervous system, there are published reports of  
27 AVMs involving all systems including the scrotum, kidney, and uterus. Herein we  
28 report a case of 37 years old male presented with recurrent gradual scrotal swelling  
29 for 4 years attributed to scrotal AVM. Embolization was done but one year later his  
30 symptoms reoccurred. As a result, left partial scrotal wall excision was carried out  
31 without complications.

32 **Keywords:** arteriovenous malformation, AVM, scrotal swelling, scrotal malformation.

33

## 34 **Introduction**

35 Arteriovenous malformations (AVMs) are benign vascular lesions. They are described  
36 as abnormal vessels fed by arteries and drained by veins without intervening  
37 capillaries.<sup>1,2</sup> Although the majority of AVMs occur in the central nervous system,  
38 there are published reports of AVMs involving the scrotum, kidney, and uterus.<sup>2,3,4</sup>  
39 Few cases of scrotal AVMs have been described in the literature. Based on the  
40 published reports, the clinical presentation of scrotal AVMs is highly variable,  
41 ranging from an incidental finding on imaging for infertility to a bleeding mass.<sup>1,5</sup>  
42 Since scrotal AVMs have variable presentations and is rarely described in the  
43 literature, we are reporting a case of a 37-year-old male presented with a slowly  
44 increasing left testicular swelling attributed to scrotal AVM.

45

## 46 **Case Report**

47 37 years old male smoker presented to the urology clinic with a gradual scrotal  
48 swelling that started four years ago. He complained of on and off scrotal pain,  
49 occasional feeling of scrotal warmth, and scrotal discomfort. The patient was  
50 diagnosed in another hospital with a testicular artery aneurysm and left testicular  
51 varicocele. The patient denied any history of trauma, urinary tract infection, voiding  
52 symptoms, previous surgeries, and his past medical history was unremarkable. Upon  
53 physical examination, the testes were intra-scrotal. There were no signs of  
54 inflammation, and the cremasteric reflex was intact bilaterally. Both epididymides  
55 were palpable and non-tender. However, pampiniform plexus at the neck of the  
56 scrotum was very pulsatile (figure A). Moreover, multiple skin varices over the left  
57 scrotum were seen. Urinalysis was normal and urine culture was negative. Routine  
58 laboratory tests were unremarkable. Abdominal and pelvis computed tomography  
59 (CT) showed left scrotal arteriovenous malformation with enlarged small and  
60 medium-sized serpiginous structures with a feeder artery arising from the proximal  
61 superficial artery. Two months later, the patient was referred to interventional  
62 radiology for embolization, which was successfully done utilizing Onyx 18% (figure  
63 B). After one year, on follow-up, the pampiniform plexus were pulsatile again which  
64 necessitated a CT angiogram. CT angiogram confirmed the recurrence of  
65 arteriovenous malformation. The patient was counselled about the available treatment  
66 options and given time to decide. Due to the risk of recurrence as well as the

67 possibility of technical failure with embolization, he decided to go with the surgical  
68 treatment. The patient was booked for surgery, and partial scrotal wall excision was  
69 done through an elliptical incision (figure C). Three arteries that feed into the  
70 arteriovenous malformation were identified and controlled with vicryl ties. The  
71 malformation and the skin that covering it were removed and sent to the pathology  
72 lab. Dartos muscle was closed in a multi-fashion layer. The skin was closed by vicryl  
73 rapide 4-0 in a vertical mattress. The histopathological study confirmed the diagnosis  
74 by detecting vascular structures extending from fibrofatty tissues measuring 8x0.5 cm  
75 grossly, and prominent subcutaneous large congested vascular spaces  
76 microscopically. The patient was discharged one day after the surgery with no  
77 complications. Two months postoperatively, the patient was doing fine with no active  
78 complaint, and the wound healed properly.

79

80 The consent was obtained orally as the images were taken from the patient in the  
81 clinic. We explained to him the importance of reporting and publishing his case for  
82 educational purposes, and he agreed.

83

#### 84 **Discussion**

85 AVMs are malformations in the circulatory system characterized by arteries and veins  
86 that are not connected by capillaries leading to various degrees of ischemia and  
87 pain.<sup>1,2</sup> Even though Central nervous system cases represent the majority of AVMs,  
88 there are published reports of AVMs involving the kidney, uterus, and scrotum.<sup>2,3,4</sup>  
89 AVMs are rarely present in the urinary tract.<sup>6</sup> We reviewed four previously published  
90 scrotal AVM cases (Table 1). All revealed ages ranging from 19 to 31 years while our  
91 patient was 37 years old. Scrotal AVM embraces wide-ranging presentations  
92 including infertility, acute recurrent pain in the hemiscrotum, pain and swelling on the  
93 testicle, and progressive diffused swelling in the scrotum with flashing skin and local  
94 warmth. Our case presented with gradually increasing left testicular swelling with on  
95 and off scrotal pain, and occasional feeling of scrotal warmth and discomfort. Of the  
96 four cases we have reviewed, three denied any history of trauma, and one had a  
97 positive trauma history which was a severe pelvic fracture, and the patient indicated  
98 that there is difficulty in maintaining erection since. Our patient denied any trauma  
99 history. Varicocele was found in two of the cases and was seen by sonography  
100 whereas our patient had multiple skin varices above the left scrotum that was seen

101 during physical examination. Each one of the four cases we reviewed diagnosed  
102 scrotal AVM with a different modality. Some were challenging and required  
103 orchiectomy for a diagnosis while others were simple and detected by pelvic  
104 angiography. Our patient was diagnosed by abdominal and pelvic CT. Two studies  
105 were able to find and embolize the feeding arteries. Our patient underwent  
106 embolization but had recurrence one-year later. Similar to our case, surgical  
107 intervention was eventually done in all four cases, and it varied from left scrotal AVM  
108 excision, orchiectomy, ileo-femoral bypass, and resection of the whole left side of the  
109 scrotum. In our case, partial scrotal wall excision was done. After the surgical  
110 intervention, all patients were symptom-free.<sup>5,6,7,8</sup> We believe that the difference in  
111 the presentation could be attributed to the location of the AVM, onset, duration, and if  
112 there is a history of trauma. A possible explanation for the differences in imaging  
113 modalities used to diagnose scrotal AVMs is the availability of imaging techniques in  
114 the hospitals that encountered those cases. The decision of surgical intervention is  
115 mainly based on the symptoms and how symptoms negatively affect the patient's  
116 quality of life.

117

### 118 **Conclusion**

119 Our case calls attention to a rare and challenging diagnosis that is scrotal AVM.  
120 Recurrent scrotal pain, swelling, and warmth together with varicocele should raise  
121 suspicion for scrotal AVM. Treatment varies depending on the symptoms present. We  
122 believe that embolization of the feeding arteries is a possible option to start with, and  
123 surgery should be preserved for recurrent cases.

124

### 125 **Authors' Contribution**

126 FMA, MSA, and SIA were responsible for conceiving the idea, literature search, data  
127 acquisition, and manuscript writing and revision. AA, SA, YA, and AB were  
128 primarily involved in the management of the case and critically reviewed the final  
129 version of the manuscript. All the authors have read and approved the final version of  
130 this manuscript.

131

132 **References**


- 133 1. Capitano M, Schieda N, Robertson SJ, et al. Images: Ruptured intratesticular  
134 arteriovenous malformation. *Can Urol Assoc J.* 2018;12(11):E489-E491.  
135 doi:10.5489/cuaj.5049
- 136 2. Lawton MT, Rutledge WC, Kim H, et al. Brain arteriovenous malformations. *Nat*  
137 *Rev Dis Primers.* 2015;1:15008. Published 2015 May 28. doi:10.1038/nrdp.2015.8
- 138 3. Wong C, Leveillee RJ, Yrizarry JM, Kirby K. Arteriovenous malformation  
139 mimicking a renal-cell carcinoma. *J Endourol.* 2002;16(9):685-686.  
140 doi:10.1089/089277902761403050
- 141 4. Szpera-Goździewicz A, Gruca-Stryjak K, Bręborowicz GH, Ropacka-Lesiak M.  
142 Uterine arteriovenous malformation - diagnosis and management. *Ginekol Pol.*  
143 2018;89(5):276-279. doi:10.5603/GP.a2018.0047
- 144 5. Monoski MA, Gonzalez RR, Thomas AJ, Goldstein M. Arteriovenous  
145 malformation of scrotum causing virtual azoospermia. *Urology.* 2006;68(1):.  
146 doi:10.1016/j.urology.2006.01.019
- 147 6. Agrawal V, Dangle P, Minhas S, Ralph D, Christopher N. Recurrent arteriovenous  
148 malformation of the scrotum secondary to pelvic trauma. *Urol Int.* 2006;77(4):365-  
149 7. doi: 10.1159/000096343. PMID: 17135789.
- 150 7. Sountoulides P, Bantis A, Asouhidou I, Aggelonidou H. Arteriovenous  
151 malformation of the spermatic cord as the cause of acute scrotal pain: a case report.  
152 *J Med Case Rep.* 2007 Oct 16;1:110. doi: 10.1186/1752-1947-1-110. PMID:  
153 17939869; PMCID: PMC2194703.
- 154 8. Mohammad A, Sahyouni W, Almerie T, Alsaid B. Angioembolization of Scrotal  
155 Arteriovenous Malformations: A Case Report and Literature Review. *Case Rep*  
156 *Vasc Med.* 2020 Feb 8;2020:8373816. doi: 10.1155/2020/8373816. PMID:  
157 32089946; PMCID: PMC7031727.



158

159 **Figure 1:** a picture of the left scrotal swelling with clear multiple skin varices.

160



161

162 **Figure 2:** the left image (A) showing a feeder artery supplying scrotal AVM. the right  
163 image (B) angiogram following Onyx embolization through the AVM is almost  
164 occluded.

165



166

167 **Figure 3:** partial left scrotal wall with AVM excision.

Author	Age	Trauma History	Presentation	Semen analysis	Varicocele	Diagnosis method	Embolization	Treatment	Follow-up
Monoski et al. <sup>5</sup>	31	No history of trauma	Infertility	Severe oligospermia	A left varicocele	Pelvic angiography	Performed	Surgical left scrotal AVM excision	3 years post-surgery, successful spontaneous pregnancy
Agrawal et al. <sup>6</sup>	25	Positive -Severe pelvic fracture 4 years ago	Pain associated with a soft swelling on his right testicle	Not Performed	No evidence of varicocele	Histopathological examination	Not performed	ileo-femoral bypass surgery	Not mentioned
Sountoulides et al. <sup>7</sup>	22	No history of trauma	Acute recurrent pain in the right hemiscrotum	Not Performed	No evidence of varicocele	Post-orchietomy specimen	Not performed	Orchiectomy	2 years post-surgery, there was no complain
Mohammed et al. <sup>8</sup>	19	No history of trauma	Progressive diffused swelling in the scrotum with flashing skin and local warmth	Not Performed	Varicocele with 1 cm dilated veins	CT arteriography	Performed	The whole left side of the scrotum was removed, and the left testicle was fixed to the right side	12 months post-surgery, there was no complain
Current case	37	No history of trauma	Gradually increasing scrotal mass with on and off scrotal pain and discomfort	Not performed	Multiple skin varices over the left scrotum were seen	Abdominal and pelvis CT	Performed	Partial scrotal wall excision	Two months post-surgery, there was no complain